arXiv:1210.4586v1 [math.FA] 16 Oct 2012

The Dirichlet heat kernel in inner uniform domains: local results, compact domains and non-symmetric forms

Janna Lierl Laurent Saloff-Coste *

October 18, 2012

Abstract

This paper provides sharp Dirichlet heat kernel estimates in inner uniform domains, including bounded inner uniform domains, in the context of certain (possibly non-symmetric) bilinear forms resembling Dirichlet forms. For instance, the results apply to the Dirichlet heat kernel associated with a uniformly elliptic divergence form operator with symmetric second order part and bounded measurable coefficients in inner uniform domains in $\mathbb{R}^n$. The results are applicable to any convex domain, to the complement of any convex domain, and to more exotic examples such as the interior and exterior of the snowflake.

AMS subject classification: 31C56,35K20,58J35,58J65,60J45,60J60

Keywords: heat equation, heat kernel, Dirichlet condition, inner uniform domains, Harnack inequality, ultracontractivity.

1 Introduction

This paper is concerned with Dirichlet heat kernel estimates in inner uniform domains. The monograph [9] introduced a general approach to this problem in the case of unbounded domains in Dirichlet spaces satisfying a global parabolic Harnack inequality. Sharp estimates for the heat kernel and the heat semigroup with Dirichlet boundary condition in domains have been studied by many authors. The article [3] contains seminal ideas. Varopoulos' work [27, 28] contains definitive results for domains above a Lipschitz function. We refer the reader to [7, 11, 17, 18, 22] for related results and further pointers to the literature. The main difference between these earlier works and the present effort is twofold. First, as in [9], our results cover inner uniform domains, a class of domains that is significantly larger than, say, Lipschitz domains. Further, inner uniformity is

---

*Both authors partially supported by NSF Grant DMS 1004771

an intrinsic notion that can be used in rather general metric spaces. This allows us to develop our results in the context of a large class of Dirichlet spaces. This larger context allows us to cover many natural and interesting examples beyond elliptic operators in $\mathbb{R}^n$, for instance, sub-elliptic operators.

This paper complements the results of [9] in several significant ways. For this purpose, we rely heavily on the results contained in the companion papers [14, 13] that were developed with the applications given here in mind.

First, we treat the case of bounded inner uniform domains which is not covered by [9]. In the unbounded case, the technique of [9] is based on using a Doob's transform that involves the "harmonic profile" h_U of the domain U, that is, a harmonic positive function in U that vanishes on the boundary (in the proper sense). In the case of bounded inner uniform domains, h_U must be replaced by the positive eigenfunction ϕ_U associated with the lowest Dirichlet eigenvalue λ_U of the domain U. This requires significant adaptation of the arguments.

Second, whether the domain is bounded or not, we include a wide class of non-symmetric second order differential operators. In the case of a fixed bounded inner uniform domain, there is not much difference in the final results between the symmetric and non symmetric cases. In the case of unbounded domains, the presence of a drift forces the estimates to be local in time (in a certain sense).

Third, in both the symmetric and non-symmetric cases, we relax the very global assumptions made in [9] to cover cases where the geometry of the underlying space is only controlled locally. In particular, we cover domains that are inner uniform only in a certain local sense. For instance, we treat the Dirichlet heat kernel for the Laplace-Beltrami operator in an unbounded inner uniform domain in a complete Riemannian manifold, without global curvature assumption or under the Ricci curvature assumption $\mathrm{Ric} \geq -\kappa g$, for some $\kappa > 0$. We also obtain some local estimates for the Dirichlet heat kernel in the interior of an unbounded convex set in $\mathbb{R}^n$. Most unbounded convex sets are not inner uniform but they are always locally uniform.

We will work in a rather abstract setting involving the notion of (not necessarily symmetric) Dirichlet forms and the associated intrinsic distance. This setting is actually very natural for this problem because, even when treating domains in $\mathbb{R}^n$, the technique we use requires the introduction of some auxiliary abstract Dirichlet spaces in which most of the work is done. Nevertheless, in the rest of this introduction, we illustrate the main results of this paper in the context of certain elliptic operators on a complete Riemannian manifold.

Let (M, g) be a complete Riemannian manifold equipped with its Riemannian measure μ. Let U be an inner uniform domain in M (for instance, if $M = \mathbb{R}^n$, bounded convex domains are inner uniform and the complement of any convex domain is inner uniform). Let L be a second order differential operator on M of the form

$$L = \Delta + X + V$$

where Δ is the Laplace-Beltrami operator on M, X is a smooth vector field on M (viewed as a differential operator acting on smooth functions $X : f \mapsto Xf = df(X)$) and V is a smooth function on M (viewed as multiplication operator). This particular structure of the differential operator L is chosen here for convenience and illustrative purpose.

Suppose that M has non-negative Ricci curvature and $X = 0, V = 0$. Suppose also that U is unbounded. Then [9] provides a global space-time two-sided estimate of the Dirichlet heat kernel $h_U^D(t, x, y)$ of the form

$$C \frac{h_U(x) h_U(y)}{\sqrt{V(x, \sqrt{t}) V(y, \sqrt{t}) h_U(x_{\sqrt{t}}) h_U(y_{\sqrt{t}})}} \exp\left(-c \frac{d_U(x, y)^2}{t}\right).$$

In this two-sided estimate, different constants $C, c \in (0, \infty)$ are used in in the lower and upper bounds. The function h_U is any fixed positive solution of $Lh = 0$ in U which vanishes at the boundary (in the proper weak sense). We call this function a harmonic profile for U. For any $x \in U$ and $r > 0$, x_r denotes a point in U with the property that $d(x, x_r) \leq Ar$ and $d(\partial U, x_r) \geq ar$ where a, A are independent of x and r. The inner uniformity of U ensures that such a point x_r exists for every $x \in U$ and $r > 0$.

One of the aims of this paper is to prove that if U is a bounded inner uniform domain in a Riemannian manifold and $L = \Delta + X + V$ as described above then the Dirichlet heat kernel for L in U can be estimated above and below on $(0, \mathrm{diam}_U^2) \times U \times U$ by

$$C \frac{e^{-t\lambda} \phi(x) \phi(y)}{\sqrt{V(x, \sqrt{t}) V(y, \sqrt{t}) \phi(x_{\sqrt{t}}) \phi(y_{\sqrt{t}})}} \exp\left(-c \frac{d_U(x, y)^2}{t}\right).$$

Here diam_U is the diameter of U, λ is the lowest Dirichlet eigenvalue of L in U and ϕ is the positive eigenfunction associated with λ and satisfying $\int \phi^2 d\mu = 1$. The notation x_r has the same meaning as before, a point in U at distance at most Ar from x and at distance at least ar from the boundary of U. The lower and upper bounds use different constants $c, C \in (0, \infty)$ and these constants depend on U and on X, V. As a corollary, we also obtain that there are constants $C = C(L, U)$ and $\alpha = \alpha(L, U)$ so that any Dirichlet eigenfunction ψ for L in U with eigenvalue λ_ψ satisfies

$$|\psi| \leq C |\lambda_\psi|^\alpha \phi.$$

In fact, we obtain a refined estimate that captures more precisely what happens at different boundary points.

For concrete examples, the reader can think of a bounded polygonal domain Ω in $\mathbb{R}^n$ as in Figure 1. In this context, we can consider the heat equation with Dirichlet boundary condition for the divergence form operator

$$Lf = \sum \partial_i^2 f + \sum b_i \partial_i f + \sum \partial_i (d_i f) + cf$$

where b_i, d_i, c are bounded measurable functions. Let ϕ be the positive eigenfunction associated with the lowest Dirichlet eigenvalue of $-L$ in Ω. Let ϕ_0

Figure 1: A polygonal domain Ω with a slit



be the positive eigenfunction associated with the lowest Dirichlet eigenvalue of $-\sum \partial_i^2$ in Ω. We show that $\phi \simeq \phi_0$ in Ω. The function ϕ_0 vanishes at different rates as x tends non-tangentially to different boundary points. The rate depends on the angle at the boundary point. For instance, ϕ will vanish linearly at smooth boundary points and will vanish quadratically when approaching the vertex of an interior right angle. The polygonal domain Ω may have a vertex with interior angle of 2π in which case the corresponding vertex is the tip of a slit. At such a vertex, ϕ vanishes as the square root of the distance to the boundary. The heat kernel estimates stated above capture this in a very precise way by reducing the estimates of the Dirichlet heat kernel to the understanding of the eigenfunction ϕ (hence ϕ_0). The case of the Koch snowflake is another good example to keep in mind.

An important special case of these results arises when the manifold M has non-negative Ricci curvature (hence satisfies the parabolic Harnack inequality at all scales) and $L = \Delta$. In this case, the results described above hold true uniformly over the class of all inner uniform domains with a specified inner uniformity constant. For instance, we can conclude that

$$\frac{e^{\lambda \mathrm{diam}_U^2} h_U^D(\mathrm{diam}_U^2, x, y)}{\phi(x)\phi(y)} \simeq 1 \text{ and } |\psi| \leq C|\lambda_\psi|^\alpha \phi.$$

uniformly over this class of domains.

The general setting in which we will work allows us to cover many different situations including the case when the Riemannian structure used above is replaced by a sub-Riemannian structure.

2 The underlying space and its geometry

2.1 The intrinsic distance

Let X be a connected, locally compact, separable metrizable space and let μ be a non-negative Borel measure on X that is finite on compact sets and positive on non-empty open sets.

We fix a symmetric, strictly local, regular Dirichlet form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F} = D(\mathcal{E}^{\mathrm{s}}))$ on $L^2(X,\mu)$ with energy measure $d\Gamma$. Recall that $d\Gamma$ is a measure-valued quadratic form defined by

$$\int f\, d\Gamma(u,u) = \mathcal{E}^{\mathrm{s}}(uf,u) - \frac{1}{2}\mathcal{E}^{\mathrm{s}}(f,u^2), \quad \forall f,u \in \mathcal{F} \cap L^\infty(X,\mu),$$

and extended to unbounded functions by setting $\Gamma(u,u) = \lim_{n\to\infty}\Gamma(u_n,u_n)$, where $u_n = \max\{\min\{u,n\},-n\}$. Using polarization, we obtain a bilinear form $d\Gamma$. In particular,

$$\mathcal{E}^{\mathrm{s}}(u,v) = \int d\Gamma(u,v), \quad \forall u,v \in \mathcal{F}.$$

We equip the Hilbert space $\mathcal{F}$ with its natural norm

$$\|f\|_{\mathcal{F}} = \left(\int_X |f|^2 d\mu + \int d\Gamma(f,f)\right)^{1/2}.$$

Let $U \subset X$ be an open set. Define

$$\mathcal{F}_{\mathrm{loc}}(U) = \{f \in L^2_{\mathrm{loc}}(U) : \forall \text{ compact } K \subset U,\ \exists f^\sharp \in \mathcal{F}, f = f^\sharp\big|_K \text{ a.e.}\}$$

For $f,g \in \mathcal{F}_{\mathrm{loc}}(U)$ we define $\Gamma(f,g)$ locally by $\Gamma(f,g)\big|_K = \Gamma(f^\sharp,g^\sharp)\big|_K$, where $K \subset U$ is compact and $f^\sharp, g^\sharp$ are functions in $\mathcal{F}$ such that $f = f^\sharp$, $g = g^\sharp$ a.e. on K. Set

$$\begin{aligned}\mathcal{F}(U) &= \{u \in \mathcal{F}_{\mathrm{loc}}(U) : \int_U |u|^2 d\mu + \int_U d\Gamma(u,u) < \infty\},\\ \mathcal{F}_{\mathrm{c}}(U) &= \{u \in \mathcal{F}(U) : \text{ the essential support of } u \text{ is compact in } U\}.\\ \mathcal{F}^0(U) &= \text{ the closure of } \mathcal{F}_{\mathrm{c}}(U) \text{ for the norm } \left(\int_U |u|^2 d\mu + \int_U d\Gamma(u,u)\right)^{1/2}.\end{aligned}$$

Definition 2.1. *The* intrinsic distance $d := d_{\mathcal{E}^{\mathrm{s}}}$ *induced by* $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ *is defined as*

$$d_{\mathcal{E}^{\mathrm{s}}}(x,y) := \sup\big\{f(x) - f(y) : f \in \mathcal{F}_{\mathrm{loc}}(X) \cap C(X),\ d\Gamma(f,f) \le d\mu\big\},$$

for all $x,y \in X$*, where* $C(X)$ *is the space of continuous functions on* X*.*

Throughout this paper, the spaces $\mathcal{F}, \mathcal{F}(U), \mathcal{F}_{\mathrm{c}}(U), \mathcal{F}^0(U)$ and the intrinsic distance d play an essential role. The space $\mathcal{F}$ is the equivalent of the Sobolev space of L^2 functions with gradient in L^2. The distance d defines the geometry of our space and will be used to introduce fundamental assumptions.

Consider the following properties of the intrinsic distance that may or may not be satisfied. They are discussed in [25, 23].

- (A1) The intrinsic distance d is finite everywhere, continuous, and defines the original topology of X.

- (A2) The space (X, d) is a complete metric space.
- (A2') $\forall x \in X, r > 0$, the open ball $B(x, r)$ is relatively compact in (X, d).

Note that if (A1) holds true then, by [25, Theorem 2], (A2) is equivalent to (A2'). Moreover, (A1)-(A2) imply that (X, d) is a geodesic space, i.e., any two points in X can be connected by a minimal geodesic in X. See [25, Theorem 1]. If (A1) and (A2) hold true then the intrinsic distance is also given by (see [23, Proposition 1])

$$d(x,y) = \sup\left\{ f(x) - f(y) : f \in \mathcal{F} \cap C_c(X),\, d\Gamma(f,f) \le d\mu \right\}, \quad x, y \in X.$$

When working in a subset Y of X, it is sometimes sufficient to assume only (A1) and

- (A2-Y) For any ball $B(x, 2r) \subset Y$, $B(x, r)$ is relatively compact.

This is a version of property (A2') that is localized in a set Y of particular interest. We will not pursue this systematically here but we will make a technical use of this fact at a later stage in the paper.

Example 2.2. *Let Ω be a domain in Euclidean space. Consider the (symmetric) Dirichlet form $\mathcal{E}_\Omega(f,f) = \int_\Omega |\nabla f|^2 d\mu$ with domain $H_0^1(\Omega)$, the Sobolev space obtained by closing the space of smooth functions with compact support in Ω in the norm $\left(\int_\Omega (|f|^2 + |\nabla f|^2) dx\right)^{1/2}$. This form is regular on Ω. The intrinsic distance is equal to the inner Euclidean distance in Ω (obtained by minimizing the length of the curves in Ω joining two points of Ω, see the next section) and property* (A1) *is satisfied. Property* (A2) *is not satisfied but* (A2-Y) *holds true for any Y with $\overline{Y} \subset \Omega$.*

2.2 Inner metric

Assume (A1)-(A2) and let Ω be a non-empty domain in X. For any continuous path $\gamma : [0,1] \to Y$, set

$$\mathrm{length}(\gamma) = \sup\left\{ \sum_{i=1}^{n} d(\gamma(t_i), \gamma(t_{i-1})) : n \in \mathbb{N}, 0 \le t_0 < \ldots < t_n \le 1 \right\}.$$

Definition 2.3. *The* inner metric *on Ω is defined as*

$$d_\Omega(x,y) = \inf\left\{ \mathrm{length}(\gamma) \middle| \gamma : [0,1] \to \Omega \text{ continuous}, \gamma(0) = x, \gamma(1) = y \right\}.$$

Let $\widetilde{\Omega}$ be the completion of Ω with respect to d_Ω.

Whenever we consider an inner ball $B_{\widetilde{\Omega}}(x,R) = \{y \in \widetilde{\Omega} : d_\Omega(x,y) < R\}$ or $B_\Omega(x,R) = B_{\widetilde{\Omega}}(x,R) \cap \Omega$, we assume that its radius is minimal in the sense that $B_{\widetilde{\Omega}}(x,R) \neq B_{\widetilde{\Omega}}(x,r)$ for all $r < R$. If x is a point in Ω, denote by $\delta(x) = \delta_\Omega(x) = d(x, \partial\Omega)$ the distance from x to the boundary of Ω. Let $\mathrm{diam}_\Omega(\Omega)$ be the diameter of Ω in the inner metric d_Ω.

Definition 2.4. *For two open sets $V \subset \Omega$, let*

$$\mathcal{F}^0_{\mathrm{loc}}(\Omega, V) = \big\{ f \in L^2_{\mathrm{loc}}(\Omega) : \forall\, W \subset V,\ \textit{rel. cpt. in } \overline{\Omega} \textit{ with } d_\Omega(W, \Omega \setminus V) > 0, \\ \exists\, f^\sharp \in \mathcal{F}^0(U) \textit{ such that } f = f^\sharp \textit{ a.e. on } W \big\}\,.$$

Definition 2.5. *Let Ω be a domain in X. For an open set $V \subset \Omega$, let $V^\sharp$ be the largest open set in $\widetilde{\Omega}$ which is contained in the closure of V in $\widetilde{\Omega}$ and whose intersection with Ω is V.*

Lemma 2.6. *Let V be an open set in Ω. A function $g \in \mathcal{F}_{\mathrm{loc}}(V)$ is in $\mathcal{F}^0_{\mathrm{loc}}(\Omega, V)$ if and only if we have $fg \in \mathcal{F}^0(\Omega)$ for any bounded function $f \in \mathcal{F}(\Omega)$ with compact support in $V^\sharp$ and such that $d\Gamma(f,f)/d\mu \in L^\infty(\Omega, \mu)$.*

Proof. See [9, Lemma 2.46]. ☐

2.3 The doubling property and Poincaré inequality

Let $Y \subset X$ and assume that the intrinsic metric d satisfies (A1)-(A2) (more generally, (A1) and (A2-Y) suffices).

Definition 2.7. *The form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies the* volume doubling property *on Y if there exists a constant $D_Y \in (0, \infty)$ such that for every ball $B(x, 2r) \subset Y$,*

$$V(x, 2r) \le D_Y\, V(x, r), \tag{VD}$$

where $V(x, r) = \mu(B(x, r))$ denotes the volume of $B(x, r)$.

Definition 2.8. *The form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies the weak* Poincaré inequality *on Y if there exists a constant $P_Y \in (0, \infty)$ such that for any ball $B(x, 2r) \subset Y$,*

$$\forall f \in D(\mathcal{E}),\ \int_{B(x,r)} |f - f_B|^2 d\mu \le P_Y\, r^2 \int_{B(x,2r)} d\Gamma(f,f), \tag{PI}$$

where $f_B = \frac{1}{V(x,r)} \int_{B(x,r)} f d\mu$ is the mean of f over $B(x, r)$.

If $Y = X$, the properties introduced in these definitions have a very global nature as they hold uniformly at all scales and locations. It is natural to introduce a more local version of these properties.

Definition 2.9. *The form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies the volume doubling property and the weak Poincaré inequality* locally *on Y if for all $x \in Y$ there is a neighborhood $Y(x)$ of x so that the volume doubling property and the Poincaré inequality hold in $Y(x)$.*

The form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies the volume doubling property and the weak Poincaré inequality up to scale R *in Y if the volume doubling property and the Poincaré inequality hold in $B(x, 2R)$ with constants independent of x, for all $x \in Y$.*

Example 2.10. *Let (M, g) be a complete Riemannian manifold and Y an open subset of M. Equipped M with its Riemannian measure and the Dirichlet form $\mathcal{E}^{\mathrm{s}}(f_1, f_2) = \int_M g(\nabla f_1, \nabla f_2) d\mu$ with its natural domain $\mathcal{F}$ (the first Sobolev space on M). In this case, the intrinsic distance on M equals the Riemannian distance.*

- *The volume doubling property and the Poincaré inequality hold locally on on Y.*
- *If Ric $\geq -\kappa g$ on the $2R$-neighborhood of Y for some fixed $\kappa > 0$ and $R > 0$ then the volume doubling property and the Poincaré inequality hold up to scale R on Y.*
- *If Ric ≥ 0 on Y then the volume doubling property and the Poincaré inequality hold on Y.*

Example 2.11. *Let G be a unimodular Lie group equipped with its Haar measure and with a family $\{X_1, \dots, X_k\}$ of left invariant vector fields that, viewed as elements of the Lie algebra, generates the Lie algebra of G (this condition is often called the Hörmander condition). Consider the Dirichlet form $\mathcal{E}^{\mathrm{s}}(f_1, f_2) = \int_G \sum_i X_i f_1 X_i f_2 d\mu$ with its natural domain $\mathcal{F}$, the space of functions in $L^2(G, \mu)$ such that, for each i, the distribution $X_i f$ can be represented by an element of $L^2(G, \mu)$. In this case, the intrinsic distance is equal to the associated sub-Riemannian distance.*

- *The volume doubling property and the Poincaré inequality hold up to scale R on G for any fixed $R > 0$.*
- *If G has polynomial volume growth (i.e., $\exists\, A, \forall\, r > 0, \quad V(e, r) \leq Cr^A$) then the volume doubling property and the Poincaré inequality hold on G.*

See, e.g., [20, Section 5.6] and [29].

2.4 Carré du champ and Lipschitz functions

Theorem 2.12. *Suppose the model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies* (A1)-(A2)*, and the volume doubling property holds locally on X. Then for any Lipschitz function f with Lipschitz constant C_L, the energy measure $d\Gamma(f, f)$ is absolutely continuous with respect to $d\mu$ and the Radon-Nikodym derivative $\Upsilon(f, f) = d\Gamma(f, f)/d\mu$ satisfies*

$$\Upsilon(f, f) \leq C_L^2$$

almost everywhere.

Proof. See [12, Theorem 2.1, Remark 2.1(ii)]. ☐

The next corollary is used to prove Proposition 6.7 and Lemma 6.10.

Corollary 2.13. *Let Ω be a domain in X. Suppose the model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies* (A1)-(A2-$\overline{\Omega}$)*, and the volume doubling property holds locally on $\overline{\Omega}$. Then any function f on Ω which is Lipschitz with respect to d_Ω with Lipschitz constant C_L is in $\mathcal{F}_{\mathrm{loc}}(\Omega)$ and satisfies*

$$C_L \geq \sup_{\Omega} \sqrt{\Upsilon(f, f)}.$$

Proof. Follows from Theorem 2.12 and a simple adaption of the arguments in [10, Corollary 3.6], [30] or [9, Corollary 2.22]. ☐

3 Inner uniformity

Let $X, \mu, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d$ be as above and assume that (A1)-(A2) are satisfied so that (X, d) is a complete metric space.

3.1 Inner uniform domains

Definition 3.1. *Fix $c \in (0,1)$, $C \in (1,\infty)$. Let Ω be a domain in X. Let $\gamma : [\alpha, \beta] \to \Omega$ be a rectifiable curve in Ω. We say that γ is a (c, C)-*uniform *curve in Ω if the following two conditions are satisfied:*

(i) $\forall t \in [\alpha, \beta]$, $\delta_\Omega\big(\gamma(t)\big) \geq c \min\big\{d\big(\gamma(\alpha), \gamma(t)\big), d\big(\gamma(t), \gamma(\beta)\big)\big\}$

(ii) $\mathrm{length}(\gamma) \leq C\, d\big(\gamma(\alpha), \gamma(\beta)\big)$.

*The domain Ω is called (c, C)-*uniform *if any two points in Ω can be joined by a (c, C)-uniform curve in Ω.*

Definition 3.2. *Fix $c \in (0,1)$, $C \in (1,\infty)$.*

*(i) Let $\gamma : [\alpha, \beta] \to \Omega$ be a rectifiable curve in Ω. We say that γ is a (c, C)-*inner uniform *curve in Ω if its is (c, C)-uniform in Ω in $(\widetilde{\Omega}, d_\Omega)$.*

(ii) We say that the domain Ω is inner uniform *if Ω is uniform in $(\widetilde{\Omega}, d_\Omega)$.*

Remark 3.3. The notions of (c, C)-length-uniformity and inner-(c, C)-length-uniformity are defined analogously by replacing $d(\gamma(s), \gamma(t))$ by $\mathrm{length}(\gamma\big|_{[s,t]})$ in condition *(i)*. The argument used in [16, Lemma 2.7] and [9, Proposition 3.3] show that if γ is a $(c$-$C)$-uniform curve in Ω joining x and y of length at most R and if the doubling property holds in $B(x, 2R)$ then there is a (c', C')-length uniform curve joining x and y in Ω. For our purpose, this means that uniformity (resp. inner uniformity) and length-uniformity (resp. inner-length-uniformity) are equivalent notions.

Lemma 3.4. *Let Ω be a (c_u, C_u)-inner uniform domain in (X, d). For every ball $B = B_{\widetilde{\Omega}}(x, r)$ in $(\widetilde{\Omega}, d_\Omega)$ with minimal radius, there exists a point $x_r \in B$ with $d_\Omega(x, x_r) = r/4$ and $d(x_r, \widetilde{\Omega} \setminus \Omega) \geq c_u r/8$.*

Proof. This is immediate, see [9, Lemma 3.20]. □

Proving that a domain Ω is inner uniform is a difficult task. In fact, we lack a general method of constructing inner uniform domains in, say, complete metric length spaces. On the other hand, many domains in Euclidean space are inner uniform.

Example 3.5. *In Euclidean space, any bounded convex domain is uniform. In addition, if Ω is convex and $B(x, aR) \subset \Omega \subset B(x, AR)$ then the uniformity constants c_u, C_u depend only on a, A. Any bounded domain with piecewise smooth boundary with a finite number of singularities and non-zero interior angle at* *each of the singularities is inner uniform. The open unit ball in* $\mathbb{R}^n$*,* $n \geq 2$*, with the trace of the half-hyperplane* $\{x : x_n = 0, x_{n-1} < 0\}$ *deleted is inner uniform. The interior and exterior of the Koch snowflake are inner uniform domains (in fact, uniform). The exterior of any convex set is inner uniform.*

Example 3.6. *Let* $G = \mathbb{R}^3$ *be the Heisenberg group with law*

$$g_1g_2 = (x_1 + x_2, y_1 + y_2, z_1 + z_2 + (1/2)(x_1y_2 - x_2y_1)), \;\; g_i = (x_i, y_i, z_i).$$

Let X *and* Y *be the left invariant vector fields on* G *with* $X(0) = \partial_x$*,* $Y(0) = \partial_y$*. Let* $\mathcal{E}(f, f) = \int_G(|Xf|^2 + |Yf|^2)d\mu$ *where* μ *denotes the Haar measure on* G *and the domain of* $\mathcal{E}$ *is the closure of smooth compactly functions for the norm* $(\int(|f|^2 + |Xf|^2 + |Yf|^2)d\mu)^{1/2}$*. Let* d *be the corresponding intrinsic distance. Examples of uniform domains include any coordinate half-space through the origin, the coordinate unit cube in* $\mathbb{R}^3$ *and any metric ball* $B(x, r)$ *in* (G, d)*. See [5, 6] and [9] for further pointers to the literature.*

3.2 Local inner uniformity

In [14], the authors derived a scale invariant boundary Harnack principle under a local version of inner uniformity which we now recall.

Definition 3.7. *Fix* $c_u \in (0, 1)$*,* $C_u \in (1, \infty)$ *and a domain* Ω*. For a point* $\xi \in \widetilde{\Omega}$*, let* $R(\Omega, \xi) \in [0, \infty]$ *be the largest* $R \geq 0$ *so that*

- *(i)* $40R/c_u \leq \operatorname{diam}_\Omega(\Omega)$ *(this is a non-trivial condition only when* Ω *is a bounded domain),*
- *(ii) Any two points in* $B_{\widetilde{\Omega}}(\xi, 20R/c_u)$ *can be connected by a curve that is* (c_u, C_u)*-inner uniform in* Ω*.*

Remark 3.8. It easily follows from Definition 3.7 that if ξ is such that $R(\Omega, \xi) > 0$ then there exists $\eta > 0$ such that

$$d_\Omega(\xi, \xi') < R(\Omega, \xi) \Longrightarrow R(\Omega, \xi') > \eta R(\Omega, \xi).$$

Consider non-empty domains $W \subset \Omega \subset X$. Let $W^\sharp$ be the largest open set in $(\widetilde{\Omega}, d_\Omega)$ whose intersection with Ω is W.

Remark 3.9. Any inner ball in $(\widetilde{W}, d_W)$ that lies in $W^\sharp$ is also an inner ball in $(\widetilde{\Omega}, d_\Omega)$. However, the metrics d_W and d_Ω do not necessarily coincide.

Definition 3.10. *Fix non-empty domains* $W \subset \Omega \subset X$*.*

- *(i) We say that* Ω *is locally inner uniform near* W *if for any point* $\xi \in W^\sharp$ *we have* $R(\Omega, \xi) > 0$ *.*
- *(ii) We say that* Ω *is locally inner uniform up to scale* $R > 0$ *near* W *if for any point* $\xi \in W^\sharp$*, we have* $R(\Omega, \xi) \geq R$*.*

Remark 3.11. (i) From these definitions, it follows easily that if Ω is (c_u, C_u)-inner uniform then $R(\Omega, \xi) \simeq \operatorname{diam}_\Omega(\Omega)$ for each $\xi \in \widetilde{\Omega}$. The constants implicitly contained in the notation $\simeq$ depend only on c_u, C_u.

(ii) By Remark 3.8, if Ω is locally inner uniform near W and $\xi \in W^\sharp$, then there exists R_ξ such that Ω is locally inner uniform up to scale R_ξ near $B_\Omega(\xi, R_\xi)$.

(iii) Assume that Ω is locally (c_u, C_u)-inner uniform up to scale R near W. Then for any point $\xi \in W^\sharp$ and $r \in (0, R)$ there exists a point $\xi_r \in \Omega$ such that $d_\Omega(\xi, \xi_r) = r/4$ and $d(\xi_r, \widetilde{\Omega} \setminus \Omega) \geq c_u r/8$. See Lemma 3.4 and [9, Lemma 3.20].

(iv) In $\mathbb{R}^n$, any domain with smooth boundary is locally inner uniform. Many such domains (e.g., an unbounded "turnip" domain) are not locally inner uniform up to scale R.

4 Adapted forms

In this section, we introduce a large class of real bilinear forms on $L^2(X, d\mu)$ that all share a common domain $\mathcal{F}$, the domain of our model form $\mathcal{E}^{\mathrm{s}}$. Further, these forms are of the type $\mathcal{E}^{\mathrm{s}} +$ lower order terms. Our goal is to pick one of these forms, $\mathcal{E}$, and to study the Dirichlet heat kernel (and Dirichlet semigroup) associated to $\mathcal{E}$ in a domain U under the hypothesis that U is inner uniform or, more generally, locally inner uniform.

4.1 First and zero order parts

Given a bilinear form $\mathcal{E}$, we set

$$\mathcal{E}^{\mathrm{sym}}(f,g) = \frac{1}{2}(\mathcal{E}(f,g) + \mathcal{E}(g,f)) \quad \text{and} \quad \mathcal{E}^{\mathrm{skew}}(f,g) = \frac{1}{2}(\mathcal{E}(f,g) - \mathcal{E}(g,f)).$$

These are, respectively, the symmetric and skew part of $\mathcal{E}$. For $f, g \in \mathcal{F}_{\mathrm{c}} \cap L^\infty(X, \mu)$, we also set

$$\mathcal{L}(f,g) = \frac{1}{2}\left(\mathcal{E}^{\mathrm{skew}}(fg, 1) + \mathcal{E}^{\mathrm{skew}}(f,g)\right) \quad \text{and} \quad \mathcal{R}(f,g) = -\mathcal{L}(g,f).$$

Obviously,

$$\mathcal{E}^{\mathrm{skew}}(f,g) = \mathcal{L}(f,g) + \mathcal{R}(f,g).$$

We recall the following definition taken from [13].

Definition 4.1. *Assuming $\mathcal{E}$ is local with $D(\mathcal{E}) = \mathcal{F}$, we say that $\mathcal{E}^{\mathrm{skew}}$ is a* first order skew form relative to $\mathcal{F}$ *if the following two properties hold:*

- *For any $u, v, f \in \mathcal{F} \cap \mathcal{C}_{\mathrm{c}}(X)$, we have*

$$\mathcal{L}(uf, v) = \mathcal{L}(u, fv) + \mathcal{L}(f, uv).$$

- *Let* $v, u_1, u_2, \ldots, u_m \in \mathcal{F} \cap \mathcal{C}_{\mathrm{c}}(X)$ *and* $u = (u_1, \ldots, u_m)$*. If* $\Phi \in C^2(\mathbb{R}^m)$*, then* $\Phi(u), \Phi_{x_i}(u) \in \mathcal{F}_{\mathrm{loc}}(X) \cap L^\infty_{\mathrm{loc}}(X, \mu)$ *and*

$$\mathcal{L}(\Phi(u), v) = \sum_{i=1}^{m} \mathcal{L}(u_i, \Phi_{x_i}(u)v).$$

Definition 4.2. *We say that the form* $(\mathcal{E}, D(\mathcal{E}))$ *is adapted to* $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ *if* $\mathcal{E}$ *is local, its domain* $D(\mathcal{E})$ *is* $\mathcal{F}$ *and:*

(i) The form $\mathcal{E}$ *satisfies*

$$\forall\, f, g \in \mathcal{F}, \;\; |\mathcal{E}(f,g)| \le C\|f\|_{\mathcal{F}}\|g\|_{\mathcal{F}},$$

and, for all $f, g \in \mathcal{F}$ *with* $fg \in \mathcal{F}_{\mathrm{c}}$*,*

$$|\mathcal{E}(fg, 1)| + |\mathcal{E}(1, fg)| \le C\|f\|_{\mathcal{F}}\|g\|_{\mathcal{F}}.$$

(ii) The symmetric bilinear form $\mathcal{E}^{\mathrm{sym}}(f,g) - \mathcal{E}^{\mathrm{sym}}(fg,1)$*, extended by continuity to* $\mathcal{F}$*, is equal to the model form* $\mathcal{E}^{\mathrm{s}}$*.*

(iii) The skew part $\mathcal{E}^{\mathrm{skew}}$ *is of first order with respect to* $\mathcal{F}$*.*

Definition 4.3. *A bilinear symmetric form* Z *is said to be a zero order form adapted to* $\mathcal{F}$ *if it is defined on* $\mathcal{F}$ *and satisfies*

$$Z(f,g) = Z(fg, 1), f, g \in \mathcal{F}, fg \in \mathcal{F}_{\mathrm{c}},$$

and

$$|Z(f,g)| \le C\|f\|_{\mathcal{F}}\|g\|_{\mathcal{F}}.$$

Since $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ is fixed throughout, we will simply say that $(\mathcal{E}, D(\mathcal{E}))$ is an adapted Dirichlet form and that Z is an adapted symmetric zero order form. Note that if $\mathcal{E}$ is an adapted form then its symmetric zero order part $Z_{\mathcal{E}}(f,g) = \mathcal{E}^{\mathrm{sym}}(fg,1)$ is a zero order form adapted to $\mathcal{F}$. Further, $\mathcal{E} = \mathcal{E}^{\mathrm{s}} + \mathcal{E}^{\mathrm{skew}} + Z_{\mathcal{E}}$.

4.2 Quantitative assumptions on the forms

We now introduce the fundamental quantitative assumptions on the bilinear forms for which we will study weak solutions of the heat equation with Dirichlet boundary condition.

Assumption A. *The form* $(\mathcal{E}, D(\mathcal{E}))$ *is a bilinear form on* $L^2(X,\mu)$ *which is adapted to the model form* $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$*. Let* $C_0 = C_0(\mathcal{E})$ *be the constant in the sector condition* $|\mathcal{E}^{\mathrm{skew}}(f,g)| \le C_0\|f\|_{\mathcal{F}}\|g\|_{\mathcal{F}}$*. Assume further that:*

(i) There are constants $C_2(\mathcal{E}), C_3(\mathcal{E}) \in [0,\infty)$ *so that for all* $f \in \mathcal{F}$ *with* $f^2 \in \mathcal{F}_{\mathrm{c}}$*,*

$$\left|\mathcal{E}^{\mathrm{sym}}(f^2,1)\right| \le 2\left(\int f^2 d\mu\right)^{\frac{1}{2}}\left(C_2(\mathcal{E})\int d\Gamma(f,f) + C_3(\mathcal{E})\int f^2 d\mu\right)^{\frac{1}{2}} \quad (1)$$

(ii) There is a constant $C_5(\mathcal{E}) \in [0,\infty)$ such that for all $f \in \mathcal{F}$, $g \in \mathcal{F}_{\mathrm{c}} \cap L^\infty(X)$,

$$\left|\mathcal{E}^{\mathrm{skew}}(f, fg^2)\right| \leq 2 \left(\int f^2 d\Gamma(g,g) \right)^{\frac{1}{2}} \left(C_5(\mathcal{E}) \int f^2 g^2 d\mu \right)^{\frac{1}{2}}. \tag{2}$$

Set

$$C_8(\mathcal{E}) := C_2(\mathcal{E}) + C_3(\mathcal{E}) + C_5(\mathcal{E}).$$

Remark 4.4. Under Assumption A the form $(\mathcal{E}, \mathcal{F})$ is closed and satisfies

$$\forall\, f \in \mathcal{F}, \quad \mathcal{E}(f,f) \geq -\alpha \|f\|_2^2,$$

with α depending only on $C_2(\mathcal{E}), C_3(\mathcal{E})$. In particular, the form $(\mathcal{E}, \mathcal{F})$ induces a continuous semigroup of bounded operators P_t on $L^2(X,\mu)$. We let $(L, D(L))$ denote the infinitesimal generator of this semigroup. By the results of [15], it is immediate that H_t is positivity preserving.

Remark 4.5. For the purpose of this work, it is essential to compare Assumption A to Assumptions 0-1-2 of [13].

(i) It is plain that any form $\mathcal{E}$ satisfying Assumption A also satisfy Assumptions 0-1-2 of [13] with respect to the model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$. Regarding Assumption 2 of [13], see [13, Remark 1.15(iv)].

(ii) Given a model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$, forms satisfying Assumption A are less general than the forms allowed by Assumptions 0-1-2 of [13]. To understand this, compare Assumption A(ii) with [13, Assumption 1(iii)] and note that Assumption A(ii) is the same as [13, Assumption 1(iii)] with $C_4 = 0$.

Remark 4.6. On Euclidean space, fix measurable bounded functions $a_{i,j}$, b_i, d_i, c, set $\mathcal{F} = D(\mathcal{E}) = W^1(\mathbb{R}^n)$ and

$$\mathcal{E}(f,g) = \int \sum_{i,j=1}^n a_{i,j} \partial_i f \partial_j g \, dx + \int \sum_{i=1}^n b_i \partial_i f \, g \, dx + \int \sum_{i=1}^n f \, d_i \partial_i g \, dx + \int c f g \, dx.$$

Set $\tilde{a}_{i,j} := (a_{i,j} + a_{j,i})/2$ and $\check{a}_{i,j} = (a_{i,j} - a_{j,i})/2$. Then the symmetric part of $\mathcal{E}$ is

$$\begin{aligned} \mathcal{E}^{\mathrm{sym}}(f,g) = & \int \sum_{i,j=1}^n \tilde{a}_{i,j} \partial_i f \partial_j g \, dx + \int \sum_{i=1}^n \frac{b_i + d_i}{2} \partial_i f \, g \, dx \\ & + \int \sum_{i=1}^n f \, \frac{b_i + d_i}{2} \partial_i g \, dx + \int c f g \, dx, \end{aligned}$$

while the skew-symmetric part of $\mathcal{E}$ is

$$\mathcal{E}^{\mathrm{skew}}(f,g) = \int \sum_{i,j=1}^n \check{a}_{i,j} \partial_i f \partial_j g \, dx + \int \sum_{i=1}^n \frac{b_i - d_i}{2} \partial_i f \, g \, dx$$

$$+\int \sum_{i=1}^{n} f\,\frac{-b_i+d_i}{2}\partial_i g\,dx.$$

The symmetric part $\mathcal{E}^{\mathrm{sym}}$ can be decomposed into its strictly local part

$$\mathcal{E}^{\mathrm{s}}(f,g) = \sum_{i,j=1}^{n} \int \tilde{a}_{i,j}\partial_i f \partial_j g\,dx$$

and its symmetric zero order part given by

$$\mathcal{E}^{\mathrm{sym}}(fg,1) = \int \sum_{i=1}^{n} \frac{b_i+d_i}{2}\partial_i(fg)\,dx + \int cfg\,dx.$$

Assume that $(\tilde{a}_{i,j})$ is uniformly elliptic and set

$$\mathcal{E}^{\mathrm{s}}(f,g) = \int \sum_{i,j=1}^{n} \tilde{a}_{i,j}\partial_i f \partial_j g\,dx, \quad f,g \in \mathcal{F}.$$

On the one hand, under these hypotheses, the form $\mathcal{E}$ satisfies [13, Assumptions 0-1-2]. On the other hand, making the hypothesis that $\mathcal{E}$ is an adapted form with respect to $(\mathcal{E}^{\mathrm{s}},\mathcal{F})$ implies that the matrix $(a_{i,j})$ is symmetric, i.e., $(a_{i,j}) = (\tilde{a}_{i,j})$.

Further, under these circumstances, the constants $C_2(\mathcal{E}), C_5(\mathcal{E})$ can be taken to be equal to 0 if $b_i = d_i = 0$ for all i (i.e., if there is no drift term). The constant $C_8(\mathcal{E})$ can be taken equal to 0 if $b_i = d_i = c = 0$.

We will need the following simple Caccioppoli-type lemma. The proof is omitted.

Lemma 4.7. *Let $(\mathcal{E},\mathcal{F})$ be a form satisfying* Assumption A. *Let $u \in \mathcal{F}_{\mathrm{loc}}$ and $\psi \in \mathcal{F}_{\mathrm{c}} \cap L^\infty(X,\mu)$. For any $k_1 > 0$, we have*

$$-\mathcal{E}^{\mathrm{s}}(u,u\psi^2) \le 4k_1 \int u^2 d\Gamma(\psi,\psi) - \left(1-\frac{1}{k_1}\right)\int \psi^2 d\Gamma(u,u).$$

Moreover, for any $k_1,k_2,k_3 > 0$,

$$\begin{aligned} -\mathcal{E}(u,u\psi^2) \quad \le \quad & (4k_1 + 2k_2C_2 + k_3)\int u^2 d\Gamma(\psi,\psi) \\ & + \left(-1+\frac{1}{k_1}+2k_2C_2\right)\int \psi^2 d\Gamma(u,u) \\ & + \left(\frac{1}{k_2}+k_2C_3+\frac{C_5}{k_3}\right)\int u^2\psi^2 d\mu \end{aligned}$$

4.3 Local weak solutions

Consider an adapted form $(\mathcal{E},\mathcal{F})$. Let V be an open set. Recall that

$$\mathcal{F}(V) = \{u \in \mathcal{F}_{\mathrm{loc}}(V) : \int_V |u|^2 d\mu + \int_V d\Gamma(u,u) < \infty\}.$$

Definition 4.8. *Let V be open and $f \in \mathcal{F}_{\mathrm{c}}(V)'$, the dual space of $\mathcal{F}_{\mathrm{c}}(V)$ (identify $L^2(X,\mu)$ with its dual space using the scalar product). A function $u : V \to \mathbb{R}$ is a* local weak solution *of the Laplace equation $-Lu = f$ in V, if*

(i) $u \in \mathcal{F}_{\mathrm{loc}}(V)$,

(ii) For any function $\phi \in \mathcal{F}_{\mathrm{c}}(V)$, $\mathcal{E}(u,\phi) = \int f\phi\, d\mu$.

For a time interval I and a Hilbert space H, let $L^2(I \to H)$ be the Hilbert space of those functions $v : I \to H$ such that

$$\|v\|_{L^2(I\to H)} = \left(\int_I \|v(t)\|_H^2\, dt\right)^{1/2} < \infty.$$

Let $\mathcal{W}^1(I \to H) \subset L^2(I \to H)$ be the Hilbert space of those functions $v : I \to H$ in $L^2(I \to H)$ whose distributional time derivative v' can be represented by functions in $L^2(I \to H)$, equipped with the norm

$$\|v\|_{\mathcal{W}^1(I\to H)} = \left(\int_I \|v(t)\|_H^2 + \|v'(t)\|_H^2\, dt\right)^{1/2} < \infty.$$

Let

$$\mathcal{F}(I \times X) = L^2(I \to \mathcal{F}) \cap \mathcal{W}^1(I \to \mathcal{F}'),$$

where $\mathcal{F}'$ denotes the dual space of $\mathcal{F}$. Let

$$\mathcal{F}_{\mathrm{loc}}(I \times V)$$

be the set of all functions $u : I \times V \to \mathbb{R}$ such that for any open interval J that is relatively compact in I, and any open subset A relatively compact in V, there exists a function $u^\sharp \in \mathcal{F}(I \times X)$ such that $u^\sharp = u$ a.e. in $J \times A$. Let

$$\mathcal{F}_{\mathrm{c}}(I \times V) = \{u \in \mathcal{F}_{\mathrm{loc}}(I \times V) : u(t,\cdot) \text{ has compact support in } V \text{ for a.e. } t \in I\}.$$

Definition 4.9. *Let I be an open interval and V an open set in X. Set $Q = I \times V$. A function $u : Q \to \mathbb{R}$ is a* local weak solution *of the heat equation $\frac{\partial}{\partial t}u = Lu$ in Q, if*

(i) $u \in \mathcal{F}_{\mathrm{loc}}(Q)$,

(ii) For any open interval J relatively compact in I,

$$\forall \phi \in \mathcal{F}_{\mathrm{c}}(Q), \quad \int_J \int_V \frac{\partial}{\partial t} u\, \phi\, d\mu\, dt + \int_J \mathcal{E}(u(t,\cdot), \phi(t,\cdot))dt = 0. \qquad (3)$$

Remark 4.10. Assuming that the intrinsic distance satisfies (A1)-(A2), an equivalent definition of a local weak solution of $\frac{\partial}{\partial t}u = Lu$ on $Q = I \times V$ is

(i) $u \in L^2(I \to D(\mathcal{E}))$,

(ii) For any open interval J relatively compact in I,

$$-\int_J \int_V \frac{\partial}{\partial t}\phi\, u\, d\mu\, dt + \int_J \mathcal{E}(u(t,\cdot),\phi(t,\cdot))dt = 0,$$

for all $\phi \in \mathcal{F}(Q)$ with compact support in $J \times V$.

See [4]. The argument uses the existence of good cut-off functions provided (A1)-(A2).

4.4 Local weak solutions with Dirichlet boundary condition along ∂U

To define weak solutions with Dirichlet boundary condition, we use Definition 2.4 where the space $\mathcal{F}^0_{\mathrm{loc}}(U,V)$ is introduced.

Definition 4.11. *Let V,U be open with $V \subset U$. A function $u : V \to \mathbb{R}$ is a* local weak solution *of the Laplace equation $-Lu = f$ in V with* Dirichlet boundary condition *along ∂U if*

(i) u is a local weak solution of $-Lu = f$ in V and

(ii) $u \in \mathcal{F}^0_{\mathrm{loc}}(U,V)$.

Next we fix an open interval I and an open set V in a domain U in X and define the notion of a local weak solution in $I \times V$ with Dirichlet boundary condition along the boundary of U. Recall that $\mathcal{F}^0(U)$ is the closure of $\mathcal{F}_{\mathrm{c}}(U)$ for the norm $(\int_U |f|^2 d\mu + \int_U d\Gamma(f,f))^{1/2}$. Define

$$\mathcal{F}^0(I \times U) = L^2(I \to \mathcal{F}^0(U)) \cap \mathcal{W}^1(I \to (\mathcal{F}^0(U))').$$

For $Q = I \times V$, define $\mathcal{F}^0_{\mathrm{loc}}(U,Q)$ to be the set of all functions $v : Q \to \mathbb{R}$ such that, for any open interval $J \subset I$ relatively compact in I and any open subset $W \subset V$ relatively compact in $\overline{U}$ with $d_U(W, U \setminus V) > 0$, there exists a function $v^\sharp$ in $\mathcal{F}^0(I \times U)$ such that $u^\sharp = u$ a.e. in $I' \times W$.

Definition 4.12. *Let I be an open interval and V an open set in X. Set $Q = I \times V$. We say that a function $u : Q \to \mathbb{R}$ is a* local weak solution *of the heat equation $\frac{\partial}{\partial t}u = Lu$ in Q with Dirichlet boundary condition along ∂U if*

(i) u is a local weak solution of the heat equation in Q and

(ii) $u \in \mathcal{F}^0_{\mathrm{loc}}(U,Q)$.

5 Harnack inequalities

Harnack inequalities play an essential and central role in the results obtained in this paper. The next two subsections discuss interior Harnack inequalities and boundary Harnack inequalities, respectively.

In this section, we consider a fixed subset Y of X. We assume that the model form $(\mathcal{E}^{\mathrm{s}},\mathcal{F})$, defined in Section 2, satisfies (A1)-(A2-Y).

5.1 Interior Harnack inequalities

For any $s \in \mathbb{R}$, $\tau > 0$, $\delta \in (0,1)$ and $B(x,2r) \subset Y$, define

$$\begin{aligned}
I &= \big(s - \tau r^2, s\big) \\
B &= B(x,r) \\
Q &= I \times B \\
Q_- &= \big(s - (3+\delta)\tau r^2/4, s - (3-\delta)\tau r^2/4\big) \times \delta B \\
Q_+ &= \big(s - (1+\delta)\tau r^2/4, s\big) \times \delta B.
\end{aligned}$$

Definition 5.1. *Let $(\mathcal{E},\mathcal{F})$ be an adapted form.*

- *We say that $(\mathcal{E},\mathcal{F})$ satisfies the* parabolic Harnack inequality *on Y if, for any $\tau > 0$, $\delta \in (0,1)$, there exists a constant $H_Y(\tau,\delta) \in (0,\infty)$ such that, for any ball $B(x,2r) \subset Y$, any $s \in \mathbb{R}$, and any positive local weak solution u of the heat equation $\frac{\partial}{\partial t}u = Lu$ in Q, the following inequality holds.*

  $$\sup_{z \in Q_-} u(z) \le H_Y \inf_{z \in Q_+} u(z) \tag{PHI}$$

  Here both the supremum and the infimum are essential, i.e., computed up to sets of measure zero.

- *We say that the parabolic Harnack inequality holds locally in Y if for each $y \in Y$ there is a neighborhood V of y in Y such that* (PHI) *holds in V (in this case, the constant H_V may indeed depend on V).*

- *We say that the parabolic Harnack inequality holds up to scale R in Y there is a constant $H_Y(R)$ such that* (PHI) *holds in any ball $B(y,2R)$, $y \in Y$, with constant $H_{B(y,2R)}$ bounded above by $H_Y(R)$.*

The parabolic Harnack inequality implies the *elliptic Harnack inequality*,

$$\sup_{z \in B(x,r)} u(z) \le H'_Y \inf_{z \in B(x,r)} u(z), \tag{EHI}$$

where u is any positive function in $\mathcal{F}_{\text{loc}}(Q)$ with $Lu = 0$ weakly in $B(x,2r)$. Recall also that (PHI) implies the Hölder continuity of local weak solutions.

The following theorem gathers fundamental known results regarding the parabolic Harnack inequality.

Theorem 5.2. *Let $X, Y, \mathcal{E}^{\text{s}}, \mathcal{F}, d, \mu$ be as in* Section 2. *Let $(\mathcal{E},\mathcal{F})$ be a form satisfying* Assumption A.

(i) The symmetric strictly local regular Dirichlet form $(\mathcal{E}^{\text{s}},\mathcal{F})$ satisfies (PHI) *on Y if and only if it satisfies the volume doubling property and the Poincaré inequality on Y.*

(ii) The symmetric strictly local regular Dirichlet form $(\mathcal{E}^{\text{s}},\mathcal{F})$ satisfies (PHI) *locally (resp. up to scale R) on Y if and only if it satisfies the volume doubling property and the Poincaré inequality locally (resp. up to scale R) on Y.*

(iii) If the model form $(\mathcal{E}^{\mathrm{s}},\mathcal{F})$ *satisfies* (PHI) *locally in* Y *then the form* $(\mathcal{E},\mathcal{F})$ *satisfies* (PHI) *locally in* Y*.*

(iv) If the model form $(\mathcal{E}^{\mathrm{s}},\mathcal{F})$ *satisfies* (PHI) *locally up to scale* $R<\infty$ *in* Y *with constant* $H(\mathcal{E}^{\mathrm{s}},R)$ *then* $(\mathcal{E},\mathcal{F})$ *satisfies* (PHI) *up to scale* $R<\infty$ *in* Y *with constant* $H(\mathcal{E},R)$ *depending only on* $H(\mathcal{E}^{\mathrm{s}},R)$*, the constants* $C_1(\mathcal{E})$*–*$C_5(\mathcal{E})$ *and an upper bound on* $C_8(\mathcal{E})R^2$*.*

Remark 5.3. The first two statements of this theorem are the Dirichlet form version of the characterization of the parabolic Harnack inequality by volume doubling and Poincaré inequality. See [8, 19, 23, 24, 26].

Statements (iii)-(iv) are variations on the key fact that the parabolic Harnack inequality for the model form $(\mathcal{E}^{\mathrm{s}},\mathcal{F})$ implies (PHI) for a wide variety of other forms in the spirit of the original work of Nash, Moser and Aronson and Serrin. The proof is contained in [13, 24, 26]. In particular, (iii)-(iv) are special cases of [13, Theorem 2.13] which covers a wider class of forms, namely, forms satisfying [13, Assumptions 0-1-2].

5.2 Boundary Harnack principle

Let $(\mathcal{E},\mathcal{F})$ be an adapted form satisfying Assumption A. Let U be a domain in X. The boundary Harnack principle is concerned with positive local weak solutions of $Lu=0$ with Dirichlet boundary condition along ∂U and their behavior near the boundary.

We will use a strong version of the boundary Harnack principle which we refer to as the geometric boundary Harnack principle.

Definition 5.4. *Let* $X,\mathcal{E}^{\mathrm{s}},\mathcal{F},d,\mu$ *be as in* Section 2. *Let* $W\subset U$ *be non-empty domains in* X*. Let* $(\mathcal{E},\mathcal{F})$ *be a form satisfying* Assumption A. *Referring to local weak solutions of* $Lu=0$ *with Dirichlet boundary condition along* ∂U *where* L *is the generator associated to* $(\mathcal{E},\mathcal{F})$*, we say that:*

(i) the geometric boundary Harnack principle *holds on* U*, if there exist constants* $a_0,A_0,A_1\in(0,\infty)$*, depending only on* U*, with the following property. Let* $\xi\in\widetilde{U}\setminus U$ *and* $r\in(0,a_0\operatorname{diam}_U(U))$*. Then for any two positive weak solutions* u *and* v *of* $Lu=0$ *in* $B_U(\xi,A_0r)$ *with Dirichlet boundary condition along* ∂U*, we have*

$$\frac{u(x)}{u(x')}\le A_1\frac{v(x)}{v(x')},\quad \forall x,x'\in B_U(\xi,r).$$

(ii) the geometric boundary Harnack principle *holds locally near* W *if, for every compact set* $K\subset W^{\sharp}\setminus W$*, there exist* $A_0(K),A_1(K)$ *and* $R(K)>0$ *such that for any* $\xi\in K$*,* $r\in(0,R(K))$ *and any two positive weak solutions* u *and* v *of* $Lu=0$ *in* $B_U(\xi,A_0(K)r)$ *with Dirichlet boundary condition along* ∂U*, we have*

$$\frac{u(x)}{u(x')}\le A_1(K)\frac{v(x)}{v(x')},\quad \forall x,x'\in B_U(\xi,r).$$

(iii) the geometric boundary Harnack principle *holds up to scale R near W if we can take $A_0(K) = A_0$, $A_1(K) = A_1$ and $R(K) = R$ in the previous statement.*

The following theorem follows immediately from the main result of [14].

Theorem 5.5. *Fix $R > 0$. Let $X, \mathcal{E}^s, \mathcal{F}, d, \mu$ be as in* Section 2*. Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A*. Let $W \subset U$ be domains in X. Assume further that:*

(i) $(\mathcal{E}, \mathcal{F})$ is a Dirichlet form.

(ii) The volume doubling property and the Poincaré inequality hold up to scale R in $\overline{W}$.

(iii) The domain U is locally (c_u, C_u)-inner uniform up to scale R near W.

Then there exist constants $a_0 \in (0, 1)$, $A_0, A_1 \in (1, \infty)$ such that for any $\xi \in W^\sharp \setminus W$, $0 < r < a_0 R$, and any two non-negative local weak solutions u, v of $Lu = 0$ in $B_U(\xi, A_0 r)$ with weak Dirichlet boundary condition along ∂U, we have

$$\frac{u(x)}{u(x')} \leq A_1 \frac{v(x)}{v(x')},$$

for all $x, x' \in B_U(\xi, r)$.

The constants a_0, A_0 depend only on the local inner uniformity constants c_u, C_u near W. The constant A_1 depends only on the inner uniformity constants c_u, C_u, an upper bound on the volume doubling constant and the Poincaré inequality constants up to scale R on $\overline{W}$, the constants $C_0(\mathcal{E})$–$C_5(\mathcal{E})$ from Assumption A *which give control over the skew-symmetric part and the killing part of the Dirichlet form $\mathcal{E}$ and an upper bound on $C_8(\mathcal{E})R^2$,*

The following theorem is a direct consequence of Theorem 5.5 and the various definitions.

Theorem 5.6. *Let $X, \mathcal{E}^s, \mathcal{F}, d, \mu$ be as in* Section 2*. Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A*. Assume further that $(\mathcal{E}, \mathcal{F})$ is a Dirichlet form. Let U be a domain in X.*

(i) Fix a domain $W \subset U$, and assume that U is locally inner uniform near W. Assume also that the volume doubling property and the Poincaré inequality hold locally in $\overline{W}$. Then the geometric boundary Harnack principle holds locally in U near W.

(ii) Fix $R \in (0, \infty]$ and a domain $W \subset U$. Assume that U is locally (c_u, C_u)-inner uniform up to scale R near W and that the volume doubling property and Poincaré inequality hold up to scale R in $\overline{W}$. Then there exists $a_0 > 0$ such that the geometric boundary Harnack principle holds locally up to scale r near W for all $r < a_0 R$ with $C_8(\mathcal{E})r^2 < \infty$.

(iii) Assume that U is inner uniform and that the volume doubling property and Poincaré inequality hold in X. Assume further that $\mathcal{E} = \mathcal{E}^s$. Then the the geometric boundary Harnack principle holds true in U.

5.3 Harmonic profiles

The main idea developed in [9] in the context of strictly local symmetric Dirichlet forms is that the Dirichlet heat kernel in a domain U can be estimated in terms of the harmonic profile h_U. In this section we recall (and extend) the notion of harmonic profiles and gather some of their key properties.

Definition 5.7. *For an open subset $U \subset X$ and under* Assumption A*, consider the bilinear form $\mathcal{E}_U^D$*

$$\mathcal{E}_U^D(f,g) = \mathcal{E}(f,g), \quad f,g \in \mathcal{F}^0(U),$$

where the domain $D(\mathcal{E}_U^D) = \mathcal{F}^0(U)$ is the closure of the space $\mathcal{F}_{\mathrm{c}}(U)$ in the norm $(\mathcal{E}^{\mathrm{s}}(f,f) + \|f\|_2)^{\frac{1}{2}}$.

Under Assumption A, the form $(\mathcal{E}_U^D, D(\mathcal{E}_U^D))$ is closed, bounded below, local and regular.

Definition 5.8. *Let $X, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d, \mu$ be as in* Section 2*. Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A *and $\mathcal{E}(f,f) \geq 0$ for all $f \in \mathcal{F}$. Let U be a domain in X. A function $h \in L^2_{\mathrm{loc}}(U)$ is called an $\mathcal{E}$-*harmonic profile *in U if it satisfies the following properties:*

- *(i) h is a weak solution of $Lu = 0$ in U;*
- *(ii) $h \in \mathcal{F}^0_{\mathrm{loc}}(U)$;*
- *(iii) $h > 0$ in U.*

*Fix a domain $W \subset U$. We say that a function h in is a (U,W)-*profile for *$\mathcal{E}$ if h is defined in W and*

- *(i) h is a weak solution of $Lu = 0$ in W;*
- *(ii) $h_U \in \mathcal{F}^0_{\mathrm{loc}}(U,W)$;*
- *(iii) $h_U > 0$ in W.*

Proposition 5.9. *Let $X, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d, \mu$ be as in* Section 2*. Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A *and which is a Dirichlet form. Fix domains $W \subset U$. Assume that the volume doubling property and Poincaré inequality hold locally on $\overline{U}$.*

- *(i) Assume that U is unbounded and inner uniform near W. Then there exists a function h which is a local weak solution of $Lh = 0$ in U and is a (U,W)-profile.*
- *(ii) If U is unbounded and locally inner uniform it admits a harmonic profile h_U.*
- *(iii) If U is bounded, inner uniform, $x_0 \in U$, and $W \subset U \setminus B_U(x_0, \epsilon)$ then the Green function $h(x) = G_U(x, x_0)$ is a (U,W)-profile.*

Proof. Note that G_U denotes the Green function in U with Dirichlet boundary condition, i.e., the Green function for the form $(\mathcal{E}^D_U, D(\mathcal{E}^D_U))$. The third statement follows immediately from [14, Lemma 3.9]. See also [9, Lemma 4.7] and [14, Lemma 3.10]. Note that this result is not correct as stated for forms satisfying Assumption A but that are not Dirichlet forms.

The idea of the proof of (i)-(ii) is to construct the profile h has a limit of normalized Green functions. The details follow the same line of reasoning as in [9, Theorem 4.16] with simple adaptations using Assumption A to take care of the fact that the form $\mathcal{E}$ is not symmetric. $\square$

Proposition 5.10. *Let $X, \mathcal{E}^s, \mathcal{F}, d, \mu$ be as in* Section 2. *Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A *and which is a Dirichlet form. Fix $R > 0$. Let $W \subset U$ be domains in X. Assume that the volume doubling property and the Poincaré inequality hold locally up to scale R on $\overline{W}$ and that U is locally inner uniform up to scale R near W. Let h be a (U, W)-profile. Then there are constants K_0, K_1 such that for any inner ball $B_U(x, r)$ with $0 < K_0 r < R$, $B_U(x, K_0 r) \subset W^\sharp$, we have*

$$\forall\, y \in B_U(x, r), \;\; h(y) \le K_1 h(x_r)$$

where x_r is any point with $d_U(x, x_r) = r/4$ and $d(x_r, X \setminus W) \ge c_u r/8$. The constants K_0 depend only on the local inner uniformity constants c_u, C_u. The constant K_1 depends only on c_u, C_u,the doubling and Poincaré constants up to scale R in $\overline{W}$, the constants C_0–C_5 which give control over the skew-symmetric part and the killing part of the Dirichlet form $\mathcal{E}$ and an upper bound on $C_8 R^2$.

Proof. Compare the ratios for h and an appropriately chosen Green function. The result follows as an immediate corollary of Theorem 5.5 and the Green function estimates obtained in [14, Lemmas 3.11-3.12]. See also [9, Theorem 4.17]. $\square$

The next two propositions are straightforward applications of Proposition 5.10. The proofs follow the same line of reasoning as in [9, Theorem 4.17] and are omitted.

Proposition 5.11 (Unbounded domains)**.** *Let $X, \mathcal{E}^s, \mathcal{F}, d, \mu$ be as in* Section 2. *Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A *and which is a Dirichlet form. Fix an unbounded domain U in X and let h be a $\mathcal{E}$ harmonic profile for U.*

(i) Assume U is locally inner uniform and that the volume doubling property and the Poincaré inequality hold locally in a neighborhood Y of $\overline{U}$ in X. Then for any compact $K \subset \widetilde{U}$ there exist $r_K, \epsilon_K > 0$ and C_K such that for any $x \in K$, $r \in (0, r_K)$, we have

$$\forall\, y \in B_U(x, r), \;\; h(y) \le C_K h(x_r)$$

where $x_r \in B_U(x, r)$ is any point with $d_U(x, x_r) = r/4$ and $d(x_r, \partial U) \ge \epsilon_K r$. Further

$$V_{h^2}(x, r) = \int_{B_U(x,r)} d\mu \simeq h(x_r)^2 V(x, r).$$

(ii) Fix $R > 0$ and assume U is locally (c_u, C_u)-inner uniform up to scale R and that the volume doubling property and the Poincaré inequality hold up to scale R in $\overline{U}$. Then there exists $a_0, A_1 \in (0, \infty)$ such that for any $x \in \widetilde{U}$, $r \in (0, a_0 R)$, we have

$$\forall\, y \in B_U(x, r), \;\; h(y) \leq A_1 h(x_r)$$

where $x_r \in B_U(x, r)$ is any point with $d_U(x, x_r) = r/4$ and $d(x_r, \partial U) \geq c_u r/8$. Further

$$V_{h^2}(x, r) = \int_{B_U(x,r)} d\mu \simeq h(x_r)^2 V(x, r).$$

The constant a_0 depends only on (c_u, C_u). The constant A_1 depends only on c_u, C_u, the volume doubling and Poincaré constant up to scale R in $\overline{U}$, the constants C_0–C_5 which give control over the skew-symmetric part and the killing part of the Dirichlet form $\mathcal{E}$ and an upper bound on $C_8 R^2$.

Proposition 5.12 (Bounded domains)**.** *Let $X, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d, \mu$ be as in* Section 2. *Let $(\mathcal{E}, \mathcal{F})$ be a form satisfying* Assumption A *and which is a Dirichlet form. Fix a bounded domain U in X. Fix $R \in (0, \frac{1}{2}\mathrm{diam}_U)$ and assume U is locally (c_u, C_u)-inner uniform at scale R and that the volume doubling property and the Poincaré inequality hold up to scale R in $\overline{U}$. Then there exist A_0, A_1 such that for any $\xi \in \widetilde{U} \setminus U$ and any $(U, B_U(\xi, R))$-profile h, we have, for all x, r such that $B_U(x, A_0 r) \subset B_U(\xi, R)$,*

$$\forall\, y \in B_U(x, r), \;\; h(y) \leq A_1 h(x_r)$$

where $x_r \in B_U(x, r)$ is any point with $d_U(x, x_r) = r/4$ and $d(x_r, \partial U) \geq c_u r/8$. Further

$$V_{h^2}(x, r) = \int_{B_U(x,r)} d\mu \simeq h(x_r)^2 V(x, r).$$

The constant A_0 depends only c_u, C_u. The constant A_1 depends only on c_u, C_u, the volume doubling and Poincaré constant up to scale R in $\overline{U}$, the constants C_0–C_5 which give control over the skew-symmetric part and the killing part of the Dirichlet form $\mathcal{E}$ and an upper bound on $C_8 R^2$.

6 The h-transform technique

Throughout this section, we let $X, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d, \mu$ be as in Section 2 and fix a form $(\mathcal{E}, \mathcal{F})$ satisfying Assumption A. We also fix a domain U in X. In addition, we fix a subdomain W of U. Recall that $W^\sharp$ is the largest open set in $\widetilde{U}$ whose intersection with U is W.

Remark 6.1. Any inner ball in $(\widetilde{W}, d_W)$ that lies in $W^\sharp$ is also an inner ball in $(\widetilde{U}, d_U)$. However, the metrics d_W and d_U do not necessarily coincide on $W^\sharp$.

We assume the model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$ satisfies (A1)-(A2-$\overline{W}$), and that the volume doubling property and the Poincaré inequality hold locally on $\overline{W}$ in X.

6.1 Some structural properties of h-transforms

Definition 6.2. *Let h be positive continuous on W. Let $H : f \in L^2(W, h^2 d\mu) \to L^2(W, d\mu)$, $f \mapsto Hf = hf$. Let $(\mathcal{E}_h^{D,W}, D(\mathcal{E}_h^{D,W}))$ be the form*

$$\mathcal{E}_h^{D,W}(f,g) = \mathcal{E}(Hf, Hg), \quad f, g \in H^{-1}(\mathcal{F}^0(W)) = D(\mathcal{E}_h^{D,W}).$$

Remark 6.3. Dropping the reference to the Dirichlet condition and W an writing $\mathcal{E}_h = \mathcal{E}_h^{D,W}$, observe that:

(i) We have

$$\mathcal{E}_h^{\mathrm{sym}}(f,g) = \mathcal{E}^{\mathrm{sym}}(hf, hg), \quad \mathcal{E}_h^{\mathrm{skew}}(f,g) = \mathcal{E}^{\mathrm{skew}}(hf, hg).$$

Further, for $f, g \in \mathcal{F}_{\mathrm{c}}(W) \cap \mathcal{C}(W)$,

$$\begin{aligned} \mathcal{E}_h^{\mathrm{s}}(f,g) &= \mathcal{E}_h^{\mathrm{sym}}(f,g) - \mathcal{E}_h^{\mathrm{sym}}(fg, 1) \\ &= \mathcal{E}^{\mathrm{s}}(hf, hg) - \mathcal{E}^{\mathrm{s}}(hfg, h) = \int h^2 d\Gamma(f,g). \end{aligned}$$

Note that $\mathcal{E}_h^{\mathrm{s}}$ must be understood as being the symmetric strictly local part of $\mathcal{E}_h$ which, in general, is not the same as the h transform $(\mathcal{E}^{\mathrm{s}})_h$ of $\mathcal{E}^{\mathrm{s}}$. The two are the same exactly when $\mathcal{E}^{\mathrm{s}}(hfg, h) = 0$ for any $f, g \in \mathcal{F}_{\mathrm{c}}(W) \cap \mathcal{C}(W)$. This is the case when h is a (U, W)-profile for $\mathcal{E}^{\mathrm{s}}$.

(ii) Recall that $\mathcal{L}_h(f,g) = \frac{1}{2}(\mathcal{E}_h^{\mathrm{skew}}(fg, 1) + \mathcal{E}_h^{\mathrm{skew}}(f,g))$. Since $\mathcal{E}$ is adapted, $\mathcal{L}$ satisfies the Leibniz rule $\mathcal{L}(uf, v) = \mathcal{L}(u, fv) + \mathcal{L}(f, uv)$, $u, v, f \in \mathcal{F}_{\mathrm{c}}(W) \cap \mathcal{C}(W)$. Hence, we obtain

$$\begin{aligned} 2\mathcal{L}_h(f,g) &= \mathcal{L}(hfg, h) + \mathcal{R}(hfg, h) + \mathcal{L}(hf, hg) + \mathcal{R}(hf, hg) \\ &= \mathcal{L}(hfg, h) - \mathcal{L}(h, hfg) + \mathcal{L}(hf, hg) - \mathcal{L}(hg, hf) \\ &= \mathcal{L}(f, gh^2) + \mathcal{L}(hg, hf) - \mathcal{L}(h, hfg) \\ &\quad + \mathcal{L}(h, hfg) + \mathcal{L}(f, h^2 g) - \mathcal{L}(hg, hf) \\ &= 2\mathcal{L}(f, h^2 g). \end{aligned}$$

This shows that $\mathcal{L}_h$ satisfies the appropriate Leibniz rule and chain rule as in Definition 4.1.

(iii) Under the additional assumption that h is a (U, W)-profile for $\mathcal{E}$, we have for $f, g \in \mathcal{F}_{\mathrm{c}}(W) \cap \mathcal{C}(W)$,

$$\mathcal{E}_h^{\mathrm{sym}}(fg, 1) = \mathcal{E}^{\mathrm{sym}}(hfg, h) = \mathcal{E}^{\mathrm{sym}}(h, hfg) = \mathcal{E}^{\mathrm{skew}}(hfg, h)$$

because $\mathcal{E}(h, hfg) = 0$ under the present condition on h, f, g.

(iv) Assume that h is an $\mathcal{E}^{\mathrm{s}}$-(U, W) profile. Then, for all $f, g \in \mathcal{F}_{\mathrm{c}}(W) \cap \mathcal{C}(W)$, we have $\mathcal{E}^{\mathrm{s}}(h, hfg) = 0$. Hence, in this case,

$$\mathcal{E}_h^{\mathrm{sym}}(fg, 1) = \mathcal{E}^{\mathrm{sym}}(h, hfg) = \mathcal{E}^{\mathrm{sym}}(h^2 fg, 1).$$

(v) Assume that $h \in \mathcal{F}_{\text{loc}}$ is positive and continuous on W and satisfies

$$\forall\, u \in \mathcal{F}_{\text{c}}(W), \;\; \mathcal{E}(f,u) = \gamma \int hu\, d\mu$$

for some $\gamma \in \mathbb{R}$. Then, for all $f, g \in \mathcal{F}_{\text{c}}(W) \cap \mathcal{C}(W)$, we have

$$\mathcal{E}_h^{\text{sym}}(fg, 1) = \mathcal{E}^{\text{sym}}(h, hfg) = \mathcal{E}^{\text{skew}}(hfg, h) + \gamma \int fgh^2 d\mu.$$

Definition 6.4. *For a fixed h, positive and continuous on W, let*

$$(\mathcal{E}^{D,W,h^2}, D(\mathcal{E}^{D,W,h^2}))$$

be the Dirichlet form on $L^2(W, h^2 d\mu)$ obtained by closing

$$\mathcal{E}^{D,W,h^2}(f,g) = \int h^2 d\Gamma(f,g), \;\; f, g \in \mathcal{F}_{\text{c}}(W).$$

Let $\mathcal{F}^h = D(\mathcal{E}^{D,W,h^2})$ be the domain of this form, that is, the closure of $\mathcal{F}_{\text{c}}(W)$ for the norm

$$\|f\|_{\mathcal{F}^h} = \mathcal{F}_W^h = \left(\int_W |f|^2 h^2 d\mu + \int_W h^2 d\Gamma(f,f) \right)^{1/2}.$$

Note that, by definition, $(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h)$ is a symmetric strictly local regular Dirichlet form on W.

Lemma 6.5. *Assume that h is continuous positive on W. Then the set*

$$H^{-1}\big(\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)\big)$$

is dense in the Hilbert space

$$D(\mathcal{E}_h^{D,W}) = H^{-1}(\mathcal{F}^0(W)), \;\; \|f\|^2_{D(\mathcal{E}_h^{D,W})} = \int_W d\Gamma(hf, hf) + \int_W h^2 |f|^2 d\mu$$

and

$$\begin{aligned} H^{-1}\big(\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)\big) &= \mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu) \\ &= \mathcal{F}_{\text{c}}(W) \cap L^\infty(W, h^2\mu). \end{aligned}$$

In particular, $\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu))$ is also dense in $D(\mathcal{E}_h^{D,W}) = H^{-1}(\mathcal{F}^0(W))$.

Proof. We follow [9, Proposition 5.7]. The set $\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)$ is dense in the Hilbert space $\mathcal{F}^0(W)$. Since H is a unitary operator between the Hilbert spaces $D(\mathcal{E}_h)$ and $\mathcal{F}^0(W)$, it follows that $H^{-1}\big(\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)\big)$ is also dense in the Hilbert space $D(\mathcal{E}_h)$. Since h, $1/h$ are both in $\mathcal{F}_{\text{loc}}(W) \cap L^\infty_{\text{loc}}(W,\mu)$, the equality

$$H^{-1}\big(\mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)\big) = \mathcal{F}_{\text{c}}(W) \cap L^\infty(W,\mu)$$

follows from the fact that $\mathcal{F}_{\text{loc}}(W) \cap L^\infty_{\text{loc}}(W,\mu)$ is an algebra. □

Lemma 6.6. *Assume that h is a (U,W)-profile for either $\mathcal{E}+\gamma\langle\cdot,\cdot\rangle$ or $\mathcal{E}^{\mathrm{s}}$. Then there exists $k\in(0,\infty)$ such that, for $f,g\in\mathcal{F}_{\mathrm{c}}(W)\cap L^\infty(W,\mu)$, we have*

(i) $\int_W d\Gamma(hf,hf)\le C\left(\mathcal{E}^{D,W,h^2}(f,f)+\int_W |f|^2h^2d\mu\right)$.

(ii) $|\mathcal{E}_h^{D,W}(f,f)|\le C\left(\mathcal{E}^{D,W,h^2}(f,f)+\int_W |f|^2h^2d\mu\right)$.

(iii) $\mathcal{E}^{D,W,h^2}(f,f)\le C\left(\mathcal{E}_h^{D,W}(f,f)+k\int_W |f|^2h^2d\mu\right)$.

In particular, $D(\mathcal{E}_h^{D,W})=D(\mathcal{E}^{D,W,h^2})=\mathcal{F}^h$.

Note that $\mathcal{F}_{\mathrm{c}}(W)\cap L^\infty(W,\mu)$ is dense in the domains of both forms $\mathcal{E}_h^{D,W}$ and $\mathcal{E}^{D,W,h^2}$ so that the last statement follows from the first two.

Proof. To simplify notation, we set $\mathcal{E}_h^{D,W}=\mathcal{E}_h$ and $\mathcal{E}^{D,W,h^2}=\mathcal{E}^{h^2}$. We give the proof when h is a (U,W)-profile for $\mathcal{E}$. The case when h is a (U,W)-profile for $\mathcal{E}+\gamma\langle\cdot,\cdot\rangle$, $\gamma\neq 0$ or $\mathcal{E}^{\mathrm{s}}$ are similar. To simplify notation, we drop the explicit reference to the Dirichlet condition and the set W. For any $f\in\mathcal{F}_{\mathrm{c}}(W)\cap L^\infty(W,\mu)$, Assumption A(i) that

$$\begin{aligned}|\mathcal{E}_h(f,f)|=|\mathcal{E}_h^{\mathrm{sym}}(hf,hf)|&\le\int d\Gamma(hf,hf)+|\mathcal{E}^{\mathrm{sym}}(h^2f^2,1)|\\&\le C\left(\int h^2d\Gamma(f,f)+\int f^2d\Gamma(h,h)+\int h^2f^2d\mu\right).\end{aligned}$$

The constant C depends only on C_2, C_3. Because $\mathcal{E}(h,hf^2)=0$, we get from Lemma 4.7 that for any $k_1,k_2,k_3>0$,

$$\begin{aligned}\left(1-\frac{1}{k_1}-2k_2C_2\right)\int f^2d\Gamma(h,h)&\le(4k_1+2k_2C_2+k_3)\int h^2d\Gamma(f,f)\\&\quad+\left(\frac{1}{k_2}+k_2C_3+\frac{C_5}{k_3}\right)\int f^2h^2d\mu.\end{aligned}$$

Hence (with a different C depending only on C_2,C_3,C_5),

$$|\mathcal{E}_h(f,f)|\le C\left(\mathcal{E}^{h^2}(f,f)+\int h^2f^2d\mu\right).$$

This proves the first desired inequality.

To prove the second statement, we use the fact that, for $f\in\mathcal{F}_{\mathrm{c}}(W)\cap L^\infty(W,\mu)$, $\mathcal{E}(h,hf^2)=0$, and Assumption A(ii) to obtain

$$\begin{aligned}\mathcal{E}^{h^2}(f,f)&=\mathcal{E}_h(f,f)-\mathcal{E}^{\mathrm{sym}}(h^2f^2,1)-\mathcal{E}^{\mathrm{s}}(h,hf^2)\\&=\mathcal{E}_h(f,f)+\mathcal{E}^{\mathrm{skew}}(h,hf^2)\\&\le\mathcal{E}_h(f,f)+k_4\int h^2d\Gamma(f,f)+\frac{C_5}{k_4}\int f^2h^2d\mu,\end{aligned}$$

where $k_4 > 0$ is arbitrary. Choosing $k_4 = 1/2$, we get

$$\mathcal{E}^{h^2}(f,f) \le 2\left(\mathcal{E}_h(f,f) + 2C_5 \int f^2 h^2 d\mu\right).$$

$\square$

Proposition 6.7. *Assume that h is continuous positive on W and belongs to $\mathcal{F}^0(U,W)$. The strictly local Dirichlet form $\big(\mathcal{E}^{D,W,h^2}, D\big(\mathcal{E}^{D,W,h^2}\big)\big)$ is regular on $(W^\sharp, h^2 d\mu)$ with core $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W)$.*

Proof. We follow the proof of [9, Proposition 5.8]. As $\big(\mathcal{E}^{D,W,h^2}, D\big(\mathcal{E}^{D,W,h^2}\big)\big)$ is regular on W, $C_{\mathrm{c}}(W) \cap D\big(\mathcal{E}^{D,W,h^2}\big)$ is dense in $D\big(\mathcal{E}^{D,W,h^2}\big)$. So we only need to show that $C_{\mathrm{c}}(W^\sharp) \cap D\big(\mathcal{E}^{D,W,h^2}\big)$ is dense in $C_{\mathrm{c}}(W^\sharp)$ in the sup norm. Consider a function $f \in \mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W)$ with Lipschitz constant k. As $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W)$ is dense in $C_{\mathrm{c}}(W^\sharp)$ in sup norm, it suffices to show that $f \in D\big(\mathcal{E}^{D,W,h^2}\big)$. In view of Lemma 6.5, it suffices to show that $fh \in \mathcal{F}^0(W)$. Since $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W) \subset \mathrm{Lip}(W, d_W)$, Corollary 2.13 implies that $f \in \mathcal{F}_{\mathrm{loc}}(W)$ and

$$\Upsilon(f,f) = \frac{d\Gamma(f,f)}{d\mu} \le k^2 \quad \text{almost everywhere on } W.$$

Since f is bounded, this shows that $f \in \mathcal{F}(W)$. Let $V \subset W$ be an open set containing $\mathrm{supp}(f) \cap W$ and relatively compact in $W^\sharp$ with the property that $\mathrm{supp}(f) \subset V^\sharp \subset W^\sharp$. Applying Lemma 2.6 with $g = h \in \mathcal{F}^0(W,V)$, we obtain that $fh \in \mathcal{F}^0(W)$. $\square$

Definition 6.8. *Assume that h is continuous positive on W and belongs to $\mathcal{F}^0(U,W)$. Recall that*

$$\mathcal{F}^h = \mathcal{F}^h_W = D(\mathcal{E}^{D,W,h^2}).$$

For an open subset $V \subset W^\sharp$, let

$$\mathcal{F}^h_{\mathrm{loc}}(V) = \{f \in L^2_{\mathrm{loc}}(V, h^2 d\mu) : \forall \text{ compact } K \subset V, \ \text{ there exists } f^\sharp \in \mathcal{F}^h \text{ so that } f = f^\sharp\big|_K \text{ a.e.}\}.$$

Similarly, define $\mathcal{F}^h(V)$ and $\mathcal{F}^h_{\mathrm{c}}(V)$ in terms of $\big(\mathcal{E}^{D,W,h^2}, D\big(\mathcal{E}^{D,W,h^2}\big)\big)$.

Remark 6.9. (i) By Proposition 6.7 and Lemma 6.5, we have

$$\mathcal{F}^h_W = H^{-1}(\mathcal{F}^0(W)) = D(\mathcal{E}_h).$$

(ii) It is now plain that the symmetric strictly local regular Dirichlet form $\big(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h_W\big)$ is the strictly local part of the symmetric part of the form $(\mathcal{E}_h, D(\mathcal{E}_h))$. In particular, for any $f \in \mathcal{F}^h_{\mathrm{c}}(W^\sharp) \cap \mathcal{C}(W^\sharp)$,

$$\mathcal{E}^{\mathrm{s}}_h(f,f) = \mathcal{E}^{\mathrm{sym}}_h(f,f) - \mathcal{E}^{\mathrm{sym}}_h(f^2,1) = \int h^2 d\Gamma(f,f) = \mathcal{E}^{D,W,h^2}(f,f).$$

(iii) The space $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_U)$ is contained in $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W)$, because for any $x, y \in W^\sharp$ it holds $d_U(x,y) \le d_W(x,y)$. In fact, both spaces are the same. To see this, observe that for any $f \in \mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_W)$ with Lipschitz constant C_W and any $x, y \in W^\sharp$ with $d_U(x,y)$ strictly less than $d_W(x,y)$ we have

$$|f(x) - f(y)| \le C\big(d_U(x, \partial W \cap U) + d_U(y, \partial W \cap U)\big) \le C d_U(x,y),$$

where

$$C = \frac{\max_{z \in W^\sharp} f(z)}{d_U(\mathrm{supp}(f), \partial W \cap U)}.$$

Hence, f is in $\mathrm{Lip}_{\mathrm{c}}(W^\sharp, d_U)$ with Lipschitz constant $C_U = \max\{C_W, C\}$.

Lemma 6.10. *Assume that h is continuous positive on W and belongs to $\mathcal{F}^0(U,W)$. The metrics d_U, d_W and $d_{\mathcal{E}^{D,W,h^2}}$ coincide on any inner ball $B = B_{\widetilde{U}}(a,r)$ with $B_{\widetilde{U}}(a,3r) \subset W^\sharp$.*

Proof. Clearly, the inner metrics d_W and d_U coincide on the ball B, since B is far away from $U \setminus W$. We follow the line of reasoning in [9, Proof of Lemma 3.32] to show that the inner metrics coincide with $d_{\mathcal{E}^{D,W,h^2}}$. Fix $y, z \in B$. Then the cut-off function

$$\rho_y(x) = \max\{d_W(y,z) - d_W(y,x), 0\},$$

is a compactly supported Lipschitz function on $(W^\sharp, d_W)$ and $\rho_y \in \mathcal{F}^h_{\mathrm{loc}}(W^\sharp) \cap C(W^\sharp)$ by Proposition 6.7. Moreover, $\Upsilon(\rho_y, \rho_y) \le 1 < \infty$ a.e. on $W^\sharp$ by Corollary 2.13. Thus,

$$d_W(y,z) = \rho_y(y) - \rho_y(z) \le d_{\mathcal{E}^{D,W,h^2}}(y,z).$$

We now show the opposite inequality. Any two points $y, z \in B \cap W$ can be connected by a curve $\gamma = \gamma_{y,z}$ in W without self-intersections. Let A_γ be an open, relatively compact subset of W that contains the curve. By [25, Theorem 3] (recall that (A1) holds on $(X, \mu, \mathcal{E}^{\mathrm{s}}, D(\mathcal{E}))$), we have

$$\begin{aligned}
\mathrm{length}(\gamma) &= \sup\{u(y) - u(z) : u \in \mathcal{F}_{\mathrm{loc}}(A_\gamma) \cap C(A_\gamma), d\Gamma(u,u) \le d\mu\} \\
&= \sup\{u(y) - u(z) : u \in \mathcal{F}^h_{\mathrm{loc}}(A_\gamma) \cap C(A_\gamma), h^2 d\Gamma(u,u) \le h^2 d\mu\} \\
&\ge d_{\mathcal{E}^{D,W,h^2}}(y,z).
\end{aligned}$$

Hence, $d_W(y,z) = \inf_\gamma L(\gamma) \ge d_{\mathcal{E}^{D,W,h^2}}(y,z)$ for all $y, z \in B \cap W$. To show that d_W and $d_{\mathcal{E}^{D,W,h^2}}$ coincide on B, approximate y and z by points in $B \cap W$. ☐

Lemma 6.11. *Assume that h is a (U,W)-profile for either $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$ or $\mathcal{E}^{\mathrm{s}}$. Then the form $\big(\mathcal{E}_h, D(\mathcal{E}_h)\big)$ satisfies* Assumption A *on $(W^\sharp, h^2 d\mu)$ with respect to $(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h_W)$. Further:*

(i) If h is a (U,W)-profile for $\mathcal{E}^{\mathrm{s}}$, then the sector condition constant $C_0(\mathcal{E}_h)$ and the constants $C_2(\mathcal{E}_h), C_3(\mathcal{E}_h), C_5(\mathcal{E}_h)$ for the form $\mathcal{E}_h$ on $(W^\sharp, h^2 d\mu)$ with respect to $(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h_W)$ are all bounded by $10(1 + C_0(\mathcal{E}) + C_8(\mathcal{E}))$.

(ii) *If h is a (U,W)-profile for $\mathcal{E}+\gamma\langle\cdot,\cdot\rangle$, then the sector condition constant $C_0(\mathcal{E}_h)$ and the constants $C_2(\mathcal{E}_h), C_3(\mathcal{E}_h), C_5(\mathcal{E}_h)$ for the form $\mathcal{E}_h$ on $(W^\sharp, h^2 d\mu)$ with respect to $(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h_W)$ are all bounded by* $10(1 + C_0(\mathcal{E}) + C_8(\mathcal{E}) + |\gamma|)$.

Proof. We only treat the case when h is a (U,W)-profile for $\mathcal{E}$ (i.e., $\gamma = 0$). The other cases are similar. We refer the reader to Remark 6.3 for various algebraic computations regarding $\mathcal{E}_h$ that are relevant to this proof. Let $f, g \in \mathcal{F}^h_W \cap \mathcal{C}_c(W)$. We have

$$\mathcal{E}_h(f,g) = \mathcal{E}^{\mathrm{s}}_h(f,g) + \mathcal{E}^{\mathrm{skew}}_h(f,g) + \mathcal{E}^{\mathrm{sym}}_h(fg,1)$$

with

$$\mathcal{E}^{\mathrm{s}}_h = \mathcal{E}^{D,W,h^2}, \;\; \mathcal{E}^{\mathrm{skew}}_h(f,g) = \mathcal{E}^{\mathrm{skew}}(hf,hg)$$

and

$$\mathcal{E}^{\mathrm{sym}}_h(fg,1) = \mathcal{E}^{\mathrm{sym}}(hfg,h) = \mathcal{E}^{\mathrm{skew}}(hfg,h)$$

where the last equality follows from the fact that $\mathcal{E}(h,hfg) = 0$ and needs to be modified appropriately when h is a profile for a form different from $\mathcal{E}$. See Remark 6.3(iii)-(iv).

Using the isometry $H : L^2(W, h^2\mu) \to L^2(W,\mu)$ and Lemma 6.6 in an obvious way, we see that

$$|\mathcal{E}^{\mathrm{skew}}(hf,hg)| \le C\|f\|_{\mathcal{F}^h}\|g\|_{\mathcal{F}^h}.$$

Next, we check that

$$|\mathcal{E}^{\mathrm{skew}}(hfg,h)| \le C\|f\|_{\mathcal{F}^h}\|g\|_{\mathcal{F}^h}.$$

By Assumption A(ii), we can find a constant k such that the symmetric bilinear form

$$(f,g) \mapsto \mathcal{E}^{\mathrm{skew}}(hfg,h) + k\left(\int_W fgh^2 d\mu + \int_W h^2 d\Gamma(f,g)\right)$$

is positive definite. Applying the Cauchy-Schwarz inequality and Assumption A(ii) yield $|\mathcal{E}^{\mathrm{skew}}(hfg,h)| \le C\|f\|_{\mathcal{F}^h}\|g\|_{\mathcal{F}^h}$ as desired. These computations also yield $|\mathcal{E}_h(fg,1)| + |\mathcal{E}_h(1,fg)| \le C\|f\|_{\mathcal{F}^h}\|g\|_{\mathcal{F}^h}$. Together with 6.3(ii), these estimates show that $\mathcal{E}_h$ is adapted to $(\mathcal{E}^{\mathrm{s}}_h, \mathcal{F}^h)$.

Next, we prove that $\mathcal{E}_h$ satisfies Assumption A(i)–(ii). Assumption A(ii) for $\mathcal{E}$ yields

$$\begin{aligned}\left|\mathcal{E}^{\mathrm{sym}}_h(f^2,1)\right| &= \left|\mathcal{E}^{\mathrm{skew}}(h,hf^2)\right| \\ &\le 2\left(\int h^2 d\Gamma(f,f)\right)^{\frac{1}{2}}\left(C_5\int f^2h^2d\mu\right)^{\frac{1}{2}}.\end{aligned}$$

This is Assumption A(i) for $\mathcal{E}_h$. Assumption A(ii) for $\mathcal{E}_h$ follows immediately from Assumption A(ii) for $\mathcal{E}$. The statements about the constants are simple bookkeeping. See Remarks 6.3(ii) and 6.9(ii). □

6.2 Properties of h-transforms in inner uniform domains

We continue to work under the hypotheses made at the beginning of Section 6.

Theorem 6.12. *Assume that*

- *The volume doubling property and the Poincaré inequality (for the model form $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$) hold locally up to scale R on $\overline{W}$.*
- *The domain U is locally (c_u, C_u)-inner uniform up to scale R near W.*

Assume that h is a (U, W)-profile for either $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$ or $\mathcal{E}^{\mathrm{s}}$. Then the following properties hold:

(i) The symmetric strictly local regular Dirichlet form $\left(\mathcal{E}^{D,W,h^2}, \mathcal{F}^h_W\right)$ satisfies property (A1) *and* (A2-B) *for any inner uniform ball $B = B_{\widetilde{U}}(a, r)$ such that $B_{\widetilde{U}}(a, 3r) \subset W^\sharp$.*

(ii) There exist constants $a_0, A_0 \in (0, \infty)$ such that, for any $r \in (0, a_0 R)$ and any inner ball $B = B_{\widetilde{U}}(a, r)$ with $B = B_{\widetilde{U}}(a, A_0 r) \subset W^\sharp$, we have

$$V_{h^2}(a, 2r) \le D(W, R) V_{h^2}(a, r)$$

and, for any $f \in \mathcal{F}^h(B)$,

$$\int_B |f - f_B|^2 h^2 d\mu \le P(W, R) r^2 \int_B h^2 d\Gamma(f, f).$$

The constants a_0, A_0 depend only c_u, C_u. The constants $D(W, R)$ and $P(W, R)$ depend only on c_u, C_u, the volume doubling and Poincaré constants up to scale R in $\overline{W}$, the constants $C_0(\mathcal{E}), C_8(\mathcal{E}), |\gamma|$ and an upper bound on $(C_8(\mathcal{E}) + |\gamma|)R^2$.

The first assertion is clear by Lemma 6.10. The proof of the second assertion is done in two stages. The first stage concerns the case when h is a profile relative to the Dirichlet form $\mathcal{E}^{\mathrm{s}}$.

Proof in the case of a $\mathcal{E}^{\mathrm{s}}$-(U, W)-profile. When h is a $\mathcal{E}^{\mathrm{s}}$-(U, W)-profile, we can apply Proposition 5.10 to obtain the asserted doubling property of the volume function V_{h^2}. Note that this very crucial step is based on the boundary Harnack principle for $\mathcal{E}^{\mathrm{s}}$ (which has only been proved so far for Dirichlet forms). Since the volume function V_{h^2} has the doubling property, the stated Poincaré inequality follows by the line of reasoning explained in [9, Theorem 3.13]. See also [9, Theorem 3.27]. One may have to change the constants a_0, A_0 when passing from the volume doubling property to the Poincaré inequality. □

Proof in the case of a (U, W)-profile for $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$. Let h still be as in the first part of the proof, that is, h is a $\mathcal{E}^{\mathrm{s}}$-(U, W)-profile. Using the result proved in stage 1 together with Lemma 6.11 and [13, Theorem 2.13], it follows that there exist a_0, A_0 such that the parabolic Harnack inequality holds for $\mathcal{E}_h + \gamma\langle\cdot h, \cdot h\rangle$ on any inner ball $B = B_{\widetilde{U}}(a, r)$, $0 < r < a_0 R$ with $B = B_{\widetilde{U}}(a, A_0 r) \subset W^\sharp$ (again, we may have to adjust the constants a_0, A_0). The constant in the Harnack inequality depends only on c_u, C_u, the volume doubling and Poincaré constant up to scale R on $\overline{W}$, $C_8(\mathcal{E})$ and upper bounds on $|\gamma|$ and $(C_8(\mathcal{E}) + |\gamma|)R^2$.

In particular, if $\hat{h}$ denotes a fixed (U, W)-profile for $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$, the Harnack inequality for $\mathcal{E}+\gamma\langle\cdot,\cdot\rangle$ applies to the weak solution $\hat{h}/h$. Adjusting a_0, A_0 again if necessary, on any inner ball $B = B_{\widetilde{U}}(a, r)$ with $B = B_{\widetilde{U}}(a, A_0 r) \subset W^\sharp$, we have

$$\forall\, x \in B_{\widetilde{U}}(a, r), \;\; \hat{h}(x) \leq A_1 \hat{h}(a_r)$$

with x_r any point in $B(a, r/4)$ with $d(x_r, \partial U) > c_u r/8$. In fact, this argument proves that the forms $\mathcal{E}$ and $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$ satisfy the geometric boundary Harnack principle up to scale R near W. Namely, this proves that Theorem 5.5 still holds true without the hypothesis (i) of Theorem 5.5. Consequently, (i) and (ii) of Theorem 5.6 hold true without assuming that $(\mathcal{E}, \mathcal{F})$ is a Dirichlet form.

In any case, the boundary control obtained above for $\hat{h}$ implies that the volume function $V_{\hat{h}^2}$ is doubling in any inner balls $B(a, r)$ as above. By the same line of reasoning outlined in the first part of the proof, the Poincaré inequality holds with respect to the measure $\hat{h}^2 d\mu$ as well. □

As a corollary that has already been used in the proof above, we have the following very useful result.

Theorem 6.13. *Assume*

- *The volume doubling property and the Poincaré inequality (for the model form* $(\mathcal{E}^{\mathrm{s}}, \mathcal{F})$*) hold locally up to scale* R *on* $\overline{W}$.
- *The domain* U *is locally* (c_u, C_u)*-inner uniform up to scale* R *near* W.

Assume that h *is a* (U, W)*-profile for either* $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$ *or* $\mathcal{E}^{\mathrm{s}}$*. Then there exist constant* a_0, A_0 *such that for any inner ball* $B = B_{\widetilde{U}}(a, r)$ *with* $r \in (0, a_0 R)$ *and* $B_{\widetilde{U}}(a, A_0 r) \subset W^\sharp$*, the parabolic Harnack inequality for* $\mathcal{E}_h$ *holds in* B *up to scale* r.

Another useful result already mentioned and used in stage 2 of the proof of Theorem 6.13 is the following extension of Theorem 5.6 to the case when $\mathcal{E}$ is not a Dirichlet form.

Theorem 6.14. *Let* $X, \mathcal{E}^{\mathrm{s}}, \mathcal{F}, d, \mu$ *be as in* Section 2*. Let* $(\mathcal{E}, \mathcal{F})$ *be a form satisfying* Assumption A*. Let* U *be a domain in* X.

(i) Fix a domain $W \subset U$*, and assume that* U *is locally inner uniform near* W*. Assume also that the volume doubling property and the Poincaré inequality hold locally in* $\overline{W}$*. Then the geometric boundary Harnack principle holds locally in* U *near* W.

(ii) Fix $R > 0$ *and a domain* $W \subset U$*. Assume that* U *is locally* (c_u, C_u)*-inner uniform up to scale* R *near* W *and that the volume doubling property and Poincaré inequality hold up to scale* R *in* $\overline{W}$*. Then there exists* $a_0 >$

0 *depending only on* c_u, C_u *such that the geometric boundary Harnack principle holds locally up to scale* $a_0 R$ *near* W *with constants depending only on* c_u, C_u*, the volume doubling and Poincaré inequality constants up to scale* R *in* $\overline{W}$*,* $C_0(\mathcal{E}), C_8(\mathcal{E})$ *and an upper bound on* $C_8(\mathcal{E})R^2$*.*

7 Estimates for the Dirichlet heat kernel

Let $(\mathcal{E}^s, \mathcal{F})$ be a model form as in Section 2 and assume that it satisfies (A1)-(A2). Let $\mathcal{E}$ be a form satisfying Assumption A. Fix a domain U and consider the bilinear form $(\mathcal{E}^D_U, \mathcal{F}^0(U))$ of Definition 5.7. In this section, we derive the main results of this paper which are two-sided estimates for the kernel of the semigroup $P^D_{U,t}$ associated with $(\mathcal{E}^D_U, \mathcal{F}^0(U))$, that is, the heat kernel for $\mathcal{E}$ with Dirichlet boundary condition along ∂U. For simplicity, let us assume that the volume doubling condition and the Poincaré inequality hold locally in U. This immediately implies that the semigroup $P^D_{U,t}$ admits a continuous positive kernel $p^D_U(t,x,y)$ in U, so that

$$P^D_{U,t} f(x) = \int p^D_U(t,x,y) f(y) dy.$$

In fact, by virtue of the local Harnack inequality, this kernel is locally Hölder continuous in $(t,x,y) \in (0,\infty) \times U \times U$.

Next, let h be a positive continuous function in U and consider the form $\left(\mathcal{E}^{D,U}_h, H^{-1}(\mathcal{F}^0(U)\right)$, where the map $H : f \mapsto hf$ is the natural isometry between $L^2(U, h^2 d\mu)$ and $L^2(U, d\mu)$ as in Definition 6.2. By construction, the form $\mathcal{E}^{D,U}_h$ induces a semigroup $P^{D,U}_{h,t} : L^2(U, h^2 d\mu) \to L^2(U, h^2 d\mu)$ given by

$$P^{D,U}_{h,t} f = h^{-1} P^D_{U,t}(hf).$$

Hence, this semigroup admits a kernel $p^{D,U}_h(t,x,y)$, $(t,x,y) \in (0,\infty) \times U \times U$ and we have

$$p^D_U(t,x,y) = h(x)h(y) p^{D,U}_h(t,x,y). \tag{4}$$

Clearly, to obtain good estimates for p^D_U, it suffices to obtain good estimates for $p^{D,U}_h$.

7.1 Local Dirichlet heat kernel estimates

In this subsection, we explain how to implement the strategy outlined above to obtain heat kernel estimates for $p^D_U(t,x,y)$ at two fixed points x and y in U (these points may, in some sense, be close to the boundary).

We fix $R_x, R_y > 0$ and make the following two basic assumptions:

(i) The volume doubling property and the Poincaré inequality hold up to scale R_x in $B(x,R)$ and up to scale R_y in $B(y,R_y)$.

(ii) The domain U is locally (c_u, C_u)-inner uniform up to scale R_x near $B_U(x,R)$ and up to scale R_y near $B_U(y,R_y)$.

Next, we pick a real γ with the property that there exists a function $h = h_\gamma$ such that h is positive continuous in U and is a $(U, B_U(x,R))$-profile and a $(U, B_U(y,R))$-profile for $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$.

The following lemma provides the existence of such a pair (γ, h_γ).

Lemma 7.1. *Let $\mathcal{E}$ be a form satisfying* Assumption A. *Let U be a domain in X. Set*

$$\lambda_U = \inf\{\mathcal{E}(f,f), f \in \mathcal{F}^0(U), \|f\|_2 = 1\}.$$

(i) If U is bounded then $-\lambda_U$ is an eigenvalue for the infinitesimal generator of $P^D_{U,t}$ and the associated normalized L^2-eigenfunction $\phi = \phi_U \in \mathcal{F}^0(U)$ is positive in U.

(ii) If U is unbounded, then there exists a function $h = h_U$ which is positive continuous in U, a local weak solution of $-Lh = \lambda_U h$ in U, and both a $(U, B_U(x,R_x))$-profile and a $(U, B_U(y,R_y))$-profile for $\mathcal{E} - \lambda_U\langle\cdot,\cdot\rangle$.

Proof. Part one follows easily from Jentzsch's Theorem (see, e.g., [21, Theorem V.6.6].

For part two, we consider a relatively compact increasing exhaustion U_n of U such that $B_U(x,R_x)$ and $B_U(y,R_y)$ are contained in U_1. For each U_n, we have an eigenfunction ϕ_{U_n} with eigenvalue λ_{U_n} in U_n given by (i). Fix a point $o \in U_1$ and consider the sequence $h_n = \phi_{U_n}/\phi_{U_n}(o)$. From the definitions and [13], it easily follows that these functions all satisfy local Harnack inequalities (with constants independent of n) in their domains and are equicontinuous. This implies that some subsequence of (h_n) converges in U to a function $h \in \mathcal{F}_{\mathrm{loc}}(U)$ which is positive and a local weak solution of $-Lh = \lambda_U h$ in U. In addition, by Theorem 6.14, the functions h_n satisfy the geometric boundary Harnack principle locally in $B_U(x,R_x)$ and $B_U(y,R_y)$, uniformly in n. This easily implies that the limit h is a $(U, B_U(x,R_x))$-profile and a $(U, B_U(y,R_y))$-profile for $\mathcal{E} - \lambda_U\langle\cdot,\cdot\rangle$. See [9, Section 4.3.2]. $\square$

Remark 7.2. (i) Recall that Assumption A implies that there exists a nonnegative real α such that $\mathcal{E}(f,f) \geq -\alpha\|f\|_2^2$ for all $f \in \mathcal{F}$, and that α is bounded above in terms of $C_2(\mathcal{E})$ and $C_3(\mathcal{E})$. Hence, $\lambda_U \geq -\alpha$.

(ii) It is not hard to modify the proof of (ii) to show that for each $\gamma \leq \lambda_U$ there exists a function h_γ which is positive continuous in U, a local weak solution of $-Lh = \gamma h$ in U, and both a $(U, B_U(x,R_x))$-profile and a $(U, B_U(y,R_y))$-profile for $\mathcal{E} - \gamma\langle\cdot,\cdot\rangle$.

Theorem 7.3. *Let $\mathcal{E}$ be a form satisfying* Assumption A. *Let U be a domain in X. Fix $x, y \in U$, $T > 0$ and γ and assume that*

(i) The volume doubling property and the Poincaré inequality hold up to scale R_x in $B(x,R_x)$ and up to scale R_y in $B(y,R_y)$.

(ii) The domain U is locally (c_u, C_u)-inner uniform up to scale R_x near $B_U(x, R_x)$ and up to scale R_y near $B_U(y, R_y)$.

(iii) There exists a function $h = h_\gamma$ such that h is positive continuous in U and both a $(U, B_U(x, R_x))$-profile and a $(U, B_U(y, R_y))$-profile for $\mathcal{E} + \gamma\langle\cdot,\cdot\rangle$.

Then for all $t \in (0, T)$, $\xi \in B_{\widetilde{U}}(x, a_0 R_x)$, $\zeta \in B_{\widetilde{U}}(y, a_0 R_y)$, we have

$$p_U^D(t, \xi, \zeta) \leq \frac{A_1 h(\xi) h(\zeta) \exp(-a_1 d_U(\xi, \zeta)^2/t)}{\sqrt{V(\xi, r_x) V(\zeta, r_y)} h(\xi_{r_x}) h(\zeta_{r_y})},$$

where $r_z = \min\{\sqrt{t}, a_0 R_z\}$ for $z = x, y$, and where $z_r \in U$ denotes a point such that $d_U(z, z_r) = r/4$ and $d(z_r, \partial U) \geq c_u r/8$, for $z = \xi, \zeta$ and $r = r_x, r_y$. Further, for all $t \in (0, T)$, $\xi \in B_{\widetilde{U}}(x, a_0 R_x)$, $\zeta \in B_{\widetilde{U}}(y, a_0 R_y)$, we have

$$p_U^D(t, \xi, \zeta) \geq \frac{a_2 h(\xi) h(\zeta) \exp(-A_2 d_U(\xi, \zeta)^2/t)}{\sqrt{V(\xi, r_x) V(\zeta, r_y)} h(\xi_{r_x}) h(\zeta_{r_y})}.$$

The constant a_0 depends only on c_u, C_u. The constant a_1 depends only on $C_0(\mathcal{E})$–$C_5(\mathcal{E})$. The constants A_1, A_2, a_2 depend only on c_u, C_u, the volume doubling and Poincaré constants up to scale R_x (resp. R_y) in $B(x, R_x)$ (resp. $B(y, R_y)$), $C_0(\mathcal{E})$-$C_5(\mathcal{E})$, $|\gamma|$ and upper bounds on $(C_8(\mathcal{E}) + |\gamma|)R_x^2$, $(C_8(\mathcal{E}) + |\gamma|)R_y^2$, TR_x^{-2} and TR_y^{-2}.

Proof. By (4) these bounds can be deduced from similar heat kernel bounds for $p_h^{D,U}$. The desired bounds for $p_h^{D,U}$ follow from classical arguments (e.g., [20, Chapter 5] and [24, 26]) based on the validity of the parabolic Harnack inequality in $B_{\widetilde{U}}(x, a_0 R_x)$ and $B_{\widetilde{U}}(y, a_0 R_y)$ which follows from Theorem 6.13. □

Remark 7.4. Theorem 7.3 holds true if we replace (iii) by the assumption that h is positive continuous in U and both a $(U, B_U(x, R_x))$-profile and a $(U, B_U(y, R_y))$-profile for $\mathcal{E}^{\mathrm{s}}$.

7.2 Dirichlet heat kernel estimates in unbounded domains

In this section we prove two-sided Dirichlet heat kernel estimates in an unbounded domain U under various hypotheses. The technique of the proof is the same as in the previous section. The form $\mathcal{E}$ is a form as in Assumption A. Our minimal assumption on the unbounded domain U is that the volume doubling property and Poincaré inequality hold locally on $\overline{U}$ and that U is locally inner uniform. By [9, 4.3.2], these minimal hypotheses imply the existence of a $\mathcal{E}^{\mathrm{s}}$-harmonic profile h in U. Since $\mathcal{E}^{\mathrm{s}}$ is our model form, it is natural to think of the $\mathcal{E}^{\mathrm{s}}$-profile h as a fundamental object. Hence, the heat kernel estimates given in this section are stated in terms of h.

The first theorem of this section provides heat kernel bounds under these minimal hypotheses. In the second theorem, these minimal hypotheses are upgraded to hypotheses that hold uniformly up to scale R for some fixed $R > 0$.

Theorem 7.5. *Let $\mathcal{E}$ be a form satisfying* Assumption A. *Let U be a locally inner uniform unbounded domain in X. Assume that the volume doubling property and the Poincaré inequality hold locally on $\overline{U}$. Let h be a $\mathcal{E}^{\mathrm{s}}$-harmonic profile in U (extended as a $\mathcal{F}$-quasi-continuous function on $\widetilde{U}$). Then the Dirichlet heat kernel p^D_U has the following properties:*

(i) The function $(t,x,y) \mapsto p^D_U(t,x,y)$ is continuous on $(0,\infty) \times U \times U$ and the function

$$(t,x,y) \mapsto \frac{p^D_U(t,x,y)}{h(x)h(y)}$$

is locally Hölder continuous in $(0,\infty) \times \widetilde{U} \times \widetilde{U}$.

(ii) There exist $c, C > 0$ such that for any pair of points $x, y \in \widetilde{U}$, there exist $r_x, r_y, A = A(x,y) \in (0,\infty)$ such that for any $t > 0$ and $\xi \in B_{\widetilde{U}}(x, r_x)$, $\zeta \in_{\widetilde{U}} (y, r_y)$, we have

$$p^D_U(t,\xi,\zeta) \le \frac{A h(\xi) h(\zeta) \exp(-c d_U(\xi,\zeta)^2/t + Ct)}{\sqrt{V(\xi,\sqrt{t_x}) V(\zeta,\sqrt{t_y})}\, h(\xi_{\sqrt{t_x}}) h(\zeta_{\sqrt{t_y}})},$$

where $t_z = \min\{t, r_z^2\}$ for $z = x, y$, and where z_r is a point in U at distance at most $r/4$ from z and at distance at least $c_u r/8$ from ∂U for $z = \xi, \zeta$.

Theorem 7.6. *Let $\mathcal{E}$ be a form satisfying* Assumption A. *Let U be an unbounded domain in X that is locally (c_u, C_u)-inner uniform up to scale R. Assume that the volume doubling property and the Poincaré inequality hold up to scale R on $\overline{U}$. Let h be a $\mathcal{E}^{\mathrm{s}}$-harmonic profile in U (extended as a $\mathcal{F}$-quasi-continuous function on $\widetilde{U}$). Then the Dirichlet heat kernel p^D_U has the following properties:*

(i) The function $(t,x,y) \mapsto p^D_U(t,x,y)$ is continuous on $(0,\infty) \times U \times U$ and there exist $\kappa, a_0 \in (0,1)$ and a constant A_1 such that for any $r \in (0, a_0 R)$, $t, t' \in (0,\infty)$, $x, x', y, y' \in \widetilde{U}$ satisfying $t \ge r^2$, $|t - t'| \le r^2/4$, $d_U(x,x') \le r$, $d(y,y') \le r$, we have

$$\left| \frac{p^D_U(t,x,y)}{h(x)h(y)} - \frac{p^D_U(t',x',y')}{h(x')h(y')} \right| \le A_1 \left(\frac{\rho}{r}\right)^{\kappa} \frac{p^D_U(t + r^2, x_r, y_r)}{h(x_r)h(y_r)}.$$

where $\rho = \sqrt{|t-t'|} + d_U(x,x') + d_U(y,y')$ and z_r is a point in U at distance at most $r/4$ from z and at distance at least $c_u r/8$ from ∂U for $z = x, y$.

(ii) There exist $c, C, a_0, A_1, a_2 A_2 > 0$ such that for any pair of points $x, y \in \widetilde{U}$ and any $t > 0$, we have

$$p^D_U(t,x,y) \le \frac{A_1 h(x) h(y) \exp(-c d_U(x,y)^2/t + Ct)}{\sqrt{V(x,\sqrt{\tau}) V(y,\sqrt{\tau})}\, h(x_{\sqrt{\tau}}) h(y_{\sqrt{\tau}})}$$

and

$$p^D_U(t,x,y) \ge \frac{a_2 h(x) h(y) \exp(-A_2 d_U(x,y)^2/t - A_2 t)}{\sqrt{V(x,\sqrt{\tau}) V(y,\sqrt{\tau})}\, h(x_{\sqrt{\tau}}) h(y_{\sqrt{\tau}})}$$

where $\tau = \min\{t, (a_0 R)^2\}$.

Proof of Theorems 7.5–7.6 (outline). The proofs of the two theorems stated above follow well established lines of reasoning. The first (and crucial) step is to use (4) and estimate the kernel $p_h^{D,U}$. Indeed, by Theorem 6.13, the associated form $\mathcal{E}_h$ satisfies a parabolic Harnack inequality. The desired bounds for $p_h^{D,U}$ follow from classical arguments (e.g., [20, Chapter 5] and [24, 26]) based on the validity of the parabolic Harnack inequality. □

Remark 7.7. In this theorem, the constant a_0 depends only on c_u, C_u. The constants c, C depend only on $C_2(\mathcal{E})$–$C_5(\mathcal{E})$. The constants κ, A_1, a_2, A_2 depend only on c_u, C_u, the volume doubling and Poincaré constant on $\overline{U}$ up to scale R, $C_0(\mathcal{E})$–$C_5(\mathcal{E})$, and on an upper bound on $C_8(\mathcal{E})R^2$.

Remark 7.8. In statement *(ii)*, the denominators can be replaced by

$$V(x, \sqrt{\tau})[h(x_{\sqrt{\tau}})]^2.$$

Note the lack of x, y symmetry of the resulting bounds. This is often useful in practice.

The following corollary of the Harnack inequality for $\mathcal{E}_h$ up to scale $a_0 R$ in $\widetilde{U}$ is also of interest. We note that if $u \in \mathcal{F}^0_{\mathrm{loc}}(U, (0,\infty) \times U)$ is a local weak solution of the heat equation for $\mathcal{E}$ in U, then $u/h \in \mathcal{F}^{h,0}_{\mathrm{loc}}(U, (0,\infty) \times U)$ is a local weak solution of the heat equation for $\mathcal{E}_h$ in U. Hence u/h satisfies the Harnack inequality up to scale $a_0 R$ in $\widetilde{U}$. This and the argument given in [20, Section 5.4.3] yields the following result.

Theorem 7.9. *Let* $\mathcal{E}$ *be a form satisfying* Assumption A. *Let* U *be an unbounded domain in* X *that is locally* (c_u, C_u)*-inner uniform up to scale* R*. Assume that the volume doubling property and the Poincaré inequality hold up to scale* R *on* $\overline{U}$*. Let* h *be a* $\mathcal{E}^{\mathrm{s}}$*-harmonic profile in* U *(extended as a* $\mathcal{F}$*-quasi-continuous function on* $\widetilde{U}$*). Let* u *be a positive local weak solution of the heat equation for* $\mathcal{E}$ *in* U *with Dirichlet boundary condition along* ∂U*. Then there exists a constant* A_1 *such that for all* $0 < s < t < \infty$ *and* $x, y \in \widetilde{U}$*, we have*

$$\frac{u(s,x)}{u(t,y)} \leq A_1 \frac{h(x)}{h(y)} \exp\left(A_1 \left(1 + \frac{t-s}{s} + \frac{t-s}{R^2} + \frac{d_U(x,y)^2}{t-s} \right) \right).$$

7.3 Dirichlet heat kernel estimates in bounded domains

This section focuses on estimates in bounded inner uniform domains and relates these results to refined intrinsic ultracontractivity estimates.

Very generally, consider a positivity preserving strongly continuous semigroup P_t acting on $L^2(U, \mu)$, where U is a bounded domain, with continuous kernel $p(t,x,y)$ such that $p(t,x,y)$ is bounded for each $t > 0$. Its adjoint P_t^* (with kernel $p^*(t,x,y) = p(t,y,x)$) has the same properties. Let λ_0 be the common bottom of the L^2-spectrum of $-L$ and $-L^*$ where L and L^* are the respective infinitesimal generators. Let ϕ_0 and ϕ_0^* be the associated positive continuous L^2-normalized eigenfunctions. Following [11], we say that the pair

(P_t, P_t^*) is intrinsically ultracontractive if for each $t > 0$ there exists a constant $c(t)$ such that

$$p(t,x,y) \leq c(t)\phi_0(x)\phi_0^*(y). \tag{5}$$

For selfadjoint semigroups, intrinsic ultracontractivity was introduced in [3]. Note that if λ_ψ is an eigenvalue for P_t with L^2-normalized eigenfunction ψ (5) implies

$$|\psi| \leq ec(1/|\lambda_\psi|)^{1/2}\phi_0 \tag{6}$$

In many interesting cases, these bounds hold with $c_t = c(1 + t^{-\nu/2})e^{-t\lambda_0}$ for some $\nu > 0$. Typically, in the literature, U is a domain in $\mathbb{R}^n$ and P_t is the semigroup associated with an elliptic second order differential operator (e.g., the Laplacian) with Dirichlet boundary condition along the boundary of U. Intrinsic ultracontractivity is then viewed as a property that depends on the regularity of the boundary of U. See, e.g., [1, 2]. In particular, it follows from [1] that the heat semigroup with Dirichlet boundary condition in any bounded inner uniform domain $U \subset \mathbb{R}^n$ is intrinsically ultracontractive with $c_t = c(1 + t^{-\nu/2})e^{-\lambda_0 t}$ for some $c = c(U), \nu = \nu(U)$. Here, we obtain the following refined results.

Theorem 7.10. *Let $\mathcal{E}$ be a form satisfying* Assumption A. *Let U be a bounded domain in X that is locally (c_u, C_u)-inner uniform up to scale R. Assume that the volume doubling property and the Poincaré inequality hold up to scale R on $\overline{U}$. Let*

$$\lambda = \lambda_U = \min\{\mathcal{E}(f,f) : f \in \mathcal{F}^0(U), \|f\|_2 = 1\},$$

and let $\phi = \phi_U$ be the associated positive L^2-normalized eigenfunction (of minus the infinitesimal generator with Dirichlet boundary condition along ∂U). Then, for all $t \in (0, R^2)$, $x, y \in \widetilde{U}$, the Dirichlet heat kernel p_U^D satisfies

$$p_U^D(t,x,y) \leq \frac{A_1\phi(x)\phi(y)e^{-cd_U(x,y)^2/t}}{\sqrt{V(x,\sqrt{t})V(y,\sqrt{t})\phi(x_{\sqrt{t}})\phi(y_{\sqrt{t}})}} \tag{7}$$

and

$$p_U^D(t,x,y) \geq \frac{a_2\phi(x)\phi(y)e^{-A_2 d_U(x,y)^2/t}}{\sqrt{V(x,\sqrt{t})V(y,\sqrt{t})\phi(x_{\sqrt{t}})\phi(y_{\sqrt{t}})}} \tag{8}$$

Further, for $t > R^2$, we have

$$a_3 \leq \frac{e^{\lambda t}p_U^D(t,x,y)}{\phi(x)\phi(y)} \leq A_3 \tag{9}$$

The constant c depends only on $C_0(\mathcal{E})$–$C_5(\mathcal{E})$. The constants $A_1, a_2, A_2, a_3, A_3 \in (0,\infty)$ depend only on c_u, C_u, the volume doubling and Poincaré constants on $\overline{U}$ up to scale R, $C_0(\mathcal{E})$–$C_5(\mathcal{E})$, and on upper bounds on $(C_8(\mathcal{E}) + |\lambda|)R^2$ and diam_U/R.

Corollary 7.11. *Referring to the notation and setting of* Theorem 7.10, *there exist a bounded continuous function w on U and a real $\omega > 0$ such that*

$$\forall\, t \geq R^2, \;\; x, y \in U, \quad \left| \frac{e^{\lambda t} p_U^D(t,x,y)}{\phi(x)\phi(y)w(y)} - 1 \right| \leq A_4 e^{-\omega t}. \tag{10}$$

Further, $a_3 \leq w \leq A_3$,

$$A_4 \leq \frac{A_3}{a_3(1 - a_3/A_3)^2} \quad \text{and} \quad \omega \geq \frac{1}{R^2} \log\left(\frac{1}{1 - a_3/A_3}\right)$$

where a_3, A_3 and R are as in Theorem 7.10.

Proof. By definition, the semigroup $K_t = e^{\lambda t} P_{\phi,t}^{D,U}$ with kernel

$$K_t(x,y) = \frac{e^{\lambda t} p_U^D(t,x,y)}{\phi(x)\phi(y)}$$

with respect to $\phi^2 d\mu$ is positivity preserving and satisfies $K_t \mathbf{1}_U = \mathbf{1}_U$. It follows that its adjoint K_t^* on $L^2(U, \phi^2 d\mu)$ admits a positive continuous eigenfunction w with eigenvalue 1. We normalize w by setting $\int w\phi^2 d\mu = 1$. Obviously, $w\phi^2 d\mu$ is then an invariant probability measure for K_t and it follows from (9) that w is bounded and bounded away from 0.

In the following computation, we think of K_t and w as Markov operators, namely,

$$f \mapsto K_t f = \int K_t(\cdot, y) f(y)\phi(y)^2 d\mu(y), \;\; f \mapsto wf = \int f w \phi^2 d\mu,$$

acting on $L^p(U, w\phi^2 d\mu)$. Note that (9) implies $a_3 \leq w \leq A_3$. Hence there exists a constant $\epsilon = a_3/A_3 > 0$ such that $K_{R^2}(x,y) \geq \epsilon w(y)$. It follows that $Q(x,y) = (1-\epsilon)^{-1}(K_{R^2}(x,y) - \epsilon w(y))$ is a Markov kernel on U with respect to $\phi^2 d\mu$ and we again denote by Q the associated operator acting on $L^p(U, w\phi^2 d\mu)$. Since $w\phi^2 d\mu$ is an invariant probability measure for Q, we have $(Q - w)^n = Q^n(I - w)$. Note also that, since $Q - w = (1-\epsilon)^{-1}(K_{R^2} - w)$,

$$\sup_{x,y}\{|Q^n(x,y)/w(y) - 1|\} = \|Q^n(I - w)\|_{1\to\infty}$$

where the right-hand side is the norm of the operator $Q^n(I-w) = Q^{n-1}(Q-w)$ from $L^1(U, w\phi^2 d\mu)$ to $L^\infty(U, w\phi^2 d\mu)$. We have

$$\|Q - w\|_{1\to\infty} \leq \epsilon^{-1}(1-\epsilon)^{-1} \;\; \text{and} \;\; \|Q^{n-1}\|_{1\to 1} \leq 1.$$

Hence, we obtain

$$\sup_{x,y}\{|Q^n(x,y)/w(y) - 1|\} \leq \epsilon^{-1}(1-\epsilon)^{-1}.$$

Since $Q^n(I - w) = (Q - w)^n = (1-\epsilon)^{-n}(K_{nR^2} - w)$, this gives

$$\sup_{x,y}\{|K_{nR^2}(x,y)/w(y) - 1|\} \leq \epsilon^{-1}(1-\epsilon)^{n-1}.$$

Since $t \mapsto \sup_{x,y}\{|K_t(x,y)/w(y) - 1|\}$ is non-increasing in t, we obtain

$$\sup_{x,y}\{|K_t(x,y)/w(y) - 1|\} \leq \epsilon^{-1}(1-\epsilon)^{-2}e^{-\omega t}, \;\; \omega = -R^{-2}\log(1-\epsilon).$$

This is exactly the desired inequality. $\square$

Remark 7.12. Let ϕ^* be the positive eigenfunction associated with the bottom eigenvalue λ for the adjoint $-L^*$ of the infinitesimal generator $-L$ of $P^D_{U,t}$. From the definitions of ϕ, ϕ^*, w, we deduce that $\phi^* = w\phi$ so that we can rewrite (10) as

$$\forall\, t \geq R^2, \;\; x, y \in U, \quad \left|\frac{e^{\lambda t}p^D_U(t,x,y)}{\phi(x)\phi^*(y)} - 1\right| \leq A_4 e^{-\omega t}. \tag{11}$$

Further, we have $c\phi \leq \phi^* \leq C\phi$ for some positive constants c, C.

Corollary 7.13. *Referring to the notation and setting of* Theorem 7.10*, there exists a constant A_5 such that, if $\psi \neq \phi$ is an $L^2(U,\mu)$-normalized eigenfunction of $-L$ with eigenvalue λ_ψ then $\eta = \lambda_\psi - \lambda \geq 1/(A_5R^2)$ and*

$$\forall\, x \in U, \;\; |\psi(x)| \leq \frac{A_5\,\phi(x)}{\sqrt{V(x, 1/\sqrt{\eta})}\phi(x_{1/\sqrt{\eta}})}. \tag{12}$$

The constant A_5 depends only on c_u, C_u, the volume doubling and Poincaré constants on $\overline{U}$ up to scale R, $C_0(\mathcal{E})$–$C_5(\mathcal{E})$, and on upper bounds on $(C_8(\mathcal{E}) + |\lambda|)R^2$ and diam_U/R.

Proof. By hypothesis, we have $P^D_{U,t}\psi = e^{-t\lambda_\psi}\psi$. Hence

$$e^{\lambda t}P^{D,U}_{\phi,t}(\psi/\phi) = e^{(\lambda-\lambda_\psi)t}(\psi/\phi).$$

The previous corollary implies that

$$\lambda_\psi - \lambda \geq \omega = 1/(A_5R^2)$$

with $A_5^{-1} = \log(1 - a_3/A_3)^{-1}$. Further, for any $x \in U$ and $t \leq R^2$, (7) yields

$$\int |p^{D,U}_\phi(t,x,y)|^2\phi(y)^2 d\mu(y) \leq \frac{A_1'}{V(x,\sqrt{t})\phi(x_{\sqrt{t}})^2}$$

where A_1' depends on the same constants as A_1 in Theorem 7.10. Because $\int \psi^2\phi^2 d\mu = 1$, it follows that

$$\begin{aligned} e^{(\lambda-\lambda_\psi)t}\frac{|\psi(x)|}{\phi(x)} &= e^{\lambda t}\left|\int p^{D,U}_\phi(t,x,y)\frac{\psi(y)}{\phi(y)}\phi(y)^2 d\mu(y)\right| \\ &\leq e^{\lambda t}\left(\int |p^{D,U}_\phi(t,x,y)|^2\phi(y)^2 d\mu(y)\right)^{1/2} \\ &\leq \frac{\sqrt{A_1'}e^{t\lambda}}{\sqrt{V(x,\sqrt{t})}\phi(x_{\sqrt{t}})} \end{aligned}$$

It now suffices to choose $t \simeq 1/(\lambda_\psi - \lambda) = 1/\eta$ (which is, indeed, of order at most R^2) to obtain

$$|\psi(x)| \leq \frac{\sqrt{A_1'} e^{|\lambda| R^2} \, \phi(x)}{\sqrt{V(x, 1/\sqrt{\eta})} \, \phi(x_{1/\sqrt{\eta}})}.$$

$\square$

Remark 7.14. In Theorem 7.10, Corollary 7.11 and Corollary 7.13, consider the special case when the volume doubling property and Poincaré inequality hold globally on $(X, (\mathcal{E}^{\mathrm{s}}, \mathcal{F}), d, \mu)$. Specialize further to the case when $\mathcal{E} = \mathcal{E}^{\mathrm{s}}$. Assume that U is a (c_u, C_u)-inner uniform domain in (X, d). Then (7)-(8)-(9) and (10)-(12) hold true with $R = \mathrm{diam}_U$ and constants $A_1, a_2, A_2, a_3, A_3, A_4, A_5$ depending only on c_u, C_u.

References

[1] R. Bañuelos, *Intrinsic ultracontractivity and eigenfunction estimates for Schrödinger operators*, J. Funct. Anal., 100 (1991), pp. 181–206.

[2] R. Bañuelos and B. Davis, *A geometrical characterization of intrinsic ultracontractivity for planar domains with boundaries given by the graphs of functions*, Indiana Univ. Math. J., 41 (1992), pp. 885–913.

[3] E. B. Davies and B. Simon, *Ultracontractivity and the heat kernel for Schrödinger operators and Dirichlet Laplacians*, J. Funct. Anal., 59 (1984), pp. 335–395.

[4] N. Eldredge and L. Saloff-Coste, *Widder's representation theorem for symmetric local Dirichlet spaces.* Submitted, http://arxiv.org/pdf/1204.1926v1.pdf.

[5] A. V. Greshnov, *On sphere geometry in the Carnot-Carathéodory metric in Carnot groups*, in Algebra, geometry, analysis and mathematical physics (Russian) (Novosibirsk, 1996), Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 1997, pp. 170–173, 191.

[6] ——, *On uniform and NTA-domains on Carnot groups*, Sibirsk. Mat. Zh., 42 (2001), pp. 1018–1035, ii.

[7] A. Grigor'yan and L. Saloff-Coste, *Dirichlet heat kernel in the exterior of a compact set*, Comm. Pure Appl. Math., 55 (2002), pp. 93–133.

[8] A. A. Grigor′yan, *The heat equation on noncompact Riemannian manifolds*, Mat. Sb., 182 (1991), pp. 55–87.

[9] P. Gyrya and L. Saloff-Coste, *Neumann and Dirichlet heat kernels in inner uniform domains*, Astérisque, (2011), pp. viii+144.

[10] F. Hirsch, *Intrinsic metrics and lipschitz functions*, J. Evol. Equ., 3 (2003), pp. 11–25.

[11] P. Kim and R. Song, *Intrinsic ultracontractivity of non-symmetric diffusion semigroups in bounded domains*, Tohoku Math. J. (2), 60 (2008), pp. 527–547.

[12] P. Koskela and Y. Zhou, *Geometry and Analysis of Dirichlet forms.* http://arxiv.org/pdf/1208.4955v1.pdf.

[13] J. Lierl and L. Saloff-Coste, *Parabolic Harnack inequality for time-dependent non-symmetric Dirichlet forms.* Submitted, http://arxiv.org/pdf/1205.6493v2.pdf.

[14] ——, *Scale-invariant boundary Harnack principle in inner uniform domains.* Submitted, http://arxiv.org/pdf/1110.2763v2.pdf.

[15] Z. M. Ma and M. Röckner, *Markov processes associated with positivity preserving coercive forms*, Canad. J. Math., 47 (1995), pp. 817–840.

[16] O. Martio, *Injectivity theorems in plane and space*, Ann. Acad. Sci. Fenn. Ser. A I Math., 4, pp. 383–401.

[17] E. M. Ouhabaz, *Analysis of heat equations on domains*, vol. 31 of London Mathematical Society Monographs Series, Princeton University Press, Princeton, NJ, 2005.

[18] E. M. Ouhabaz and F.-Y. Wang, *Sharp estimates for intrinsic ultracontractivity on $C^{1,\alpha}$-domains*, Manuscripta Math., 122 (2007), pp. 229–244.

[19] L. Saloff-Coste, *A note on Poincaré, Sobolev, and Harnack inequalities*, Internat. Math. Res. Notices, (1992), pp. 27–38.

[20] L. Saloff-Coste, *Aspects of Sobolev-type inequalities*, vol. 289 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2002.

[21] H. H. Schaefer, *Banach lattices and positive operators*, Springer-Verlag, New York, 1974. Die Grundlehren der mathematischen Wissenschaften, Band 215.

[22] R. Song, *Estimates on the Dirichlet heat kernel of domains above the graphs of bounded $C^{1,1}$ functions*, Glas. Mat. Ser. III, 39(59) (2004), pp. 273–286.

[23] K.-T. Sturm, *Analysis on local Dirichlet spaces. I. Recurrence, conservativeness and L^p-Liouville properties*, J. Reine Angew. Math., 456 (1994), pp. 173–196.

[24] ———, *Analysis on local Dirichlet spaces. II. Upper Gaussian estimates for the fundamental solutions of parabolic equations*, Osaka J. Math., 32 (1995), pp. 275–312.

[25] ———, *On the geometry defined by Dirichlet forms*, vol. 36 of Progr. Probab., Birkhäuser, Basel, 1995.

[26] ———, *Analysis on local Dirichlet spaces. III. The parabolic Harnack inequality*, J. Math. Pures Appl. (9), 75 (1996), pp. 273–297.

[27] N. T. Varopoulos, *Potential theory in Lipschitz domains*, Canad. J. Math., 53 (2001), pp. 1057–1120.

[28] ———, *Gaussian estimates in Lipschitz domains*, Canad. J. Math., 55 (2003), pp. 401–431.

[29] N. T. Varopoulos, L. Saloff-Coste, and T. Coulhon, *Analysis and geometry on groups*, vol. 100 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 1992.

[30] N. Weaver, *Lipschitz algebras and derivations. ii. exterior differentiation*, J. Funct. Anal., 178 (2000), pp. 64–112.